

UC-NRLF



B 4 248 869

MATHEMATICAL MONOGRAPHS.

EDITED BY

Mansfield Merriman and Robert S. Woodward.

Octavo, Cloth, \$1.00 each.

No. 1. HISTORY OF MODERN MATHEMATICS.

By DAVID EUGENE SMITH.

No. 2. SYNTHETIC PROJECTIVE GEOMETRY.

By GEORGE BRUCE HALSTED.

No. 3. DETERMINANTS.

By LAENAS GIFFORD WELD.

No. 4. HYPERBOLIC FUNCTIONS.

By JAMES McMAHON.

No. 5. HARMONIC FUNCTIONS.

By WILLIAM E. BYERLY.

No. 6. GRASSMANN'S SPACE ANALYSIS.

By EDWARD W. HYDE.

No. 7. PROBABILITY AND THEORY OF ERRORS.

By ROBERT S. WOODWARD.

No. 8. VECTOR ANALYSIS AND QUATERNIONS.

By ALEXANDER MACFARLANE.

No. 9. DIFFERENTIAL EQUATIONS.

By WILLIAM WOOLSEY JOHNSON.

No. 10. THE SOLUTION OF EQUATIONS.

By MANSFIELD MERRIMAN.

No. 11. FUNCTIONS OF A COMPLEX VARIABLE.

By THOMAS S. FISKE.

PUBLISHED BY

JOHN WILEY & SONS, NEW YORK.

CHAPMAN & HALL, Limited, LONDON.

MATHEMATICAL MONOGRAPHS.

EDITED BY

MANSFIELD MERRIMAN AND ROBERT S. WOODWARD.

No. 6.

GRASSMANN'S
SPACE ANALYSIS.

BY

EDWARD W. HYDE,

ACTUARY OF THE COLUMBIA INSURANCE COMPANY.

FOURTH EDITION.

FIRST THOUSAND.



NEW YORK:

JOHN WILEY & SONS.

LONDON: CHAPMAN & HALL, LIMITED.

1906.

32-179

COPYRIGHT, 1896,
BY
MANSFIELD MERRIMAN AND ROBERT S. WOODWARD
UNDER THE TITLE
HIGHER MATHEMATICS.

First Edition, September, 1896.

Second Edition, January, 1898.

Third Edition, August, 1900.

Fourth Edition, January, 1906.

EDITORS' PREFACE.

THE volume called *Higher Mathematics*, the first edition of which was published in 1896, contained eleven chapters by eleven authors, each chapter being independent of the others, but all supposing the reader to have at least a mathematical training equivalent to that given in classical and engineering colleges. The publication of that volume is now discontinued and the chapters are issued in separate form. In these reissues it will generally be found that the monographs are enlarged by additional articles or appendices which either amplify the former presentation or record recent advances. This plan of publication has been arranged in order to meet the demand of teachers and the convenience of classes, but it is also thought that it may prove advantageous to readers in special lines of mathematical literature.

It is the intention of the publishers and editors to add other monographs to the series from time to time, if the call for the same seems to warrant it. Among the topics which are under consideration are those of elliptic functions, the theory of numbers, the group theory, the calculus of variations, and non-Euclidean geometry; possibly also monographs on branches of astronomy, mechanics, and mathematical physics may be included. It is the hope of the editors that this form of publication may tend to promote mathematical study and research over a wider field than that which the former volume has occupied.

December, 1905.

iii

AUTHOR'S PREFACE.

THIS little book is an attempt to present simply and concisely the elementary principles of the "Extensive Analysis" as fully developed in the comprehensive treatises of Hermann Grassmann, restricting the treatment however to the geometry of two and three dimensional space. It is designed to set forth, as far as is possible in so brief a work, the remarkable adaptability and effectiveness of the methods used as applied to the various problems and operations of geometry and mechanics.

The ideas of direction and position appear to the writer to be as simple and fundamental as that of magnitude, and an algebra which deals directly with all three of these ideas should not be greatly more difficult than the ordinary one, which deals with magnitude only. The result of using such an algebra is an extraordinary gain in the brevity of operations and the expressiveness of formulas and equations.

Some of the terms belonging to this general subject are frequently employed in modern text-books on mechanics and physics, even when no use is made of the algebraic systems from which they are derived.

It is hoped that this little book may do something to interest students, and to help toward bringing in the time when the methods as well as the ideas of this calculus shall come into general use.

CINCINNATI, O., December, 1905.

CONTENTS.

ART. 1. EXPLANATIONS AND DEFINITIONS	Page 8
2. SUM AND DIFFERENCE OF TWO POINTS	9
3. SUM OF TWO WEIGHTED POINTS	12
4. SUM OF ANY NUMBER OF POINTS	15
5. REFERENCE SYSTEMS	20
6. NATURE OF GOEMETRIC MULTIPLICATION	24
7. PLANIMETRIC PRODUCTS	26
8. THE COMPLEMENT	33
9. EQUATIONS OF CONDITION AND FORMULAS	39
10. STEREOMETRIC PRODUCTS	44
11. THE COMPLEMENT IN SOLID SPACE	50
12. ADDITION OF SECTS IN SOLID SPACE	53

GRASSMANN'S SPACE ANALYSIS.

INTRODUCTION.

The title adopted for this brief and elementary discussion of Grassmann's methods indicates at once its limitations; for his theory in its fullness treats of the linear relation

$$p = \sum x_k e_k, \quad (k = 0, 1, 2, \dots, n),$$

in which n may be any positive whole number, $x_0, x_1 \dots$ any numbers whatever, and $e_0, e_1 \dots$ units of any kind which are susceptible of being related to each other by such an equation as the above. Our treatment extends only to the case when n does not exceed three, and $e_0, e_1 \dots$ are points, or point products.

Grassmann's first publication of his new system was made in 1844 in a book entitled "*Die Lineale Ausdehnungslehre Ein Neuer Zweig der Mathematik.*" His novel and fruitful ideas were however presented in a somewhat abstruse and unusual form, with the result, as the author himself states in the preface to the second edition issued in 1878, that scarcely any notice was taken of the book by Mathematicians.

He was finally convinced that it would be necessary to treat the subject in an entirely different manner in order to gain the attention of the mathematical world. Accordingly in 1862 he published "*Die Ausdehnungslehre, vollständig und in strenger Form bearbeitet,*" in which the treatment is algebraic, and is developed from the equation given above.

Since that time his great work has been more fully appreciated, but not even yet, in the opinion of the writer, at its real value.

Hamilton first gained the ear of the English-speaking world for his Quaternion methods, and was fortunate in having some very zealous adherents and interpreters who made propaganda for them, and were inclined to undervalue work not originating in England.

It is hoped that the following brief presentation of Grassmann's Analysis will serve to interest some to the extent that they may be led to investigate his original works.

The writer has followed in the main the notation of Grassmann himself, the principal deviations being the omission of the brackets from geometric products, writing pq instead of $[pq]$, and a somewhat different treatment of the product $p|q$.

ART. 1. EXPLANATIONS AND DEFINITIONS.

The algebra with which the student is already familiar deals directly with only one quality of the various geometric and mechanical entities, such as lines, forces, etc., namely, with their magnitude. Such questions as How much? How far? How long? are answered by an algebraic operation or series of operations. Questions of direction and position are dealt with indirectly by means of systems of coordinates of various kinds. In this chapter an algebra* will be developed which deals directly with the three qualities of geometric and mechanical quantities, viz., magnitude, position, and direction. A geometric quantity may possess one, two, or all three of these properties simultaneously; thus a straight line of given length has all three, while a point has only one.

The geometric quantities with which we are to be concerned are the point, the straight line, the plane, the vector, and the plane-vector.

When the word "line" is used by itself, a "straight line" will be always intended. A portion of a given straight line of definite length will be called a "sect"; though when the length

* The algebra of this chapter is a particular case of the very general and comprehensive theory developed by Hermann Grassmann, and published by him in 1844 under the title "Die lineale Ausdehnungslehre, ein neuer Zweig der Mathematik." He published also a second treatise on the subject in 1862.

of the sect is a matter of indifference, the word line will frequently be used instead. Similarly, a definite area of a given plane will be called a "plane-sect."

If a point recede to infinity, it has no longer any significance as regards position, but still indicates a direction, since all lines passing through finite points, and also through this point at infinity, are parallel. Similarly, a line wholly at infinity fixes a plane direction, that is, all planes passing through finite points, and also through this line at infinity, are parallel. Thus a point and line at infinity are respectively equivalent to a line direction and a plane direction.

A quantity possessing magnitude only will be termed a "scalar" quantity. Such are the ordinary subjects of algebraic analysis, a , x , $\sin \theta$, $\log z$, etc., and they may evidently be intrinsically either positive or negative.

The letter T prefixed to a letter denoting some geometric quantity will be used to designate its absolute or numerical magnitude, always positive. Thus, if L be a sect, and P a plane-sect, then TL is the length of L , and TP is the area of P . That portion of a geometric quantity whose magnitude is unity will be called its "unit," and will be indicated by prefixing the letter U ; thus $UL = \text{unit of } L = \text{sect one unit long on line } L$.* Hence we have $TL \cdot UL = L$.

ART. 2. SUM AND DIFFERENCE OF TWO POINTS.

In geometric addition and subtraction we shall use the ordinary symbols $+$, $-$, $=$, but with modified significance, as will appear in the development of the subject.

Every mathematical, or other, theory rests on certain fundamental assumptions, the justification for these assumptions

* The word "scalar" and the use of the letters T and U , as above, were introduced by Hamilton in his Quaternions. T stands for tensor, i.e., stretcher, and TL is the factor that stretches UL into L . The notation $|L|$ for absolute magnitude is not used, because the sign $|$ has been appropriated by Grassmann to another use.

lying in the harmony and reasonableness of the resulting theory, and its accordance with the ascertained facts of nature.

Our first assumption, then, will be that the associative and commutative laws hold for geometric addition and subtraction, that is, whatever A , B , C may represent, we have

$$\begin{aligned} A + B + C &= (A + B) + C = A + (B + C) \\ &= A + C + B = (A + C) + B. \end{aligned}$$

We shall also assume that we always have $A - A = 0$, and that the same quantity may be added to or subtracted from both sides of an equation without affecting the equality.

Now let p_1, p_2 be two points, and consider the equation

$$p_2 + p_1 - p_1 = p_2 + (p_1 - p_1) = p_2. \quad (1)$$

In this form we have an identity. Write it, however, in the form

$$p_2 - p_1 + p_1 = (p_2 - p_1) + p_1 = p_2, \quad (2)$$

and it appears that $p_2 - p_1$ is an operator that changes p_1 into p_2 by being added to it. Conceive this change of p_1 into p_2 to take place along the straight line through p_1 and p_2 ; then the operation is that of moving a point through a definite length or distance in a definite direction, namely, from p_1 to p_2 . This operator has been called by Hamilton "a vector,"* that is, a carrier, because it carries p_1 rectilinearly to p_2 . Grassmann gives to it the name *Strecke*, and some writers now use the word "stroke" in the same sense.

Again, $p_2 - p_1$ is the difference of two points, and the only difference that can exist between them is that of position, i.e. a certain distance in a certain direction.

Hence we may regard $p_2 - p_1$ as a directed length, and also as the operator which moves p_1 over this length in this direction. Writing $p_2 - p_1 = \epsilon$, equation (2) becomes

$$p_1 + \epsilon = p_2. \quad (3)$$

* See the first of Hamilton's *Lectures on Quaternions*, where a very full discussion of equation (2) will be found. Also Grassmann (1862), Art. 227.

Thus the sum of a point and a vector is a point distant from the first by the length of the vector and in its direction.

Since $p_2 - p_1 = -(p_1 - p_2)$, it appears that the negative of a vector is a vector of the same length in the opposite direction.


If $p_2 - p_1 = 0$, or $p_2 = p_1$, p_2 must coincide with p_1 because there is now no difference between the two points.

The question arises as to what, if any, effect the operator $p_2 - p_1$ should have on any other point p_3 , that is, what is the value of the expression $p_2 - p_1 + p_3$?

We will assume that it is some point p_4 , so that we have $p_2 - p_1 + p_3 = p_4$,

$$\text{or} \quad p_2 - p_1 = p_4 - p_3. \quad (4)$$

This implies that the transference from p_3 to p_4 is the same in amount and direction as that from p_1 to p_2 , that is, that p_1, p_2, p_4, p_3 are the four corners of a parallelogram taken in order. Thus equal vectors have the same length and direction, and, conversely, vectors having the same length and direction are equal.



Note that parallel vectors of equal length are not necessarily equal, for their directions may be opposite.

Equation (4) may also be written

$$p_2 + p_3 = p_4 + p_1, \quad (5)$$

so that, whatever meaning may be assigned to the sum of two points, if we are to be consistent with assumptions already made, we must have the sum of either pair of opposite corner-points of a parallelogram equal to the sum of the other pair. The sum cannot therefore depend on the actual distances apart of the points forming the pairs, for the ratio of these two distances may be made as large or as small as we please.

If n be a scalar quantity, $n\epsilon$ will denote that the operation ϵ is to be performed n times on a point to which $n\epsilon$ is added, that is, the point will be moved n times the length of ϵ ; hence

$n\epsilon$ is a vector n times as long as ϵ , and having the same or the opposite direction according to the sign of n .

In the figure above, let

$$p_2 - p_1 = \epsilon_1, \quad p_3 - p_1 = \epsilon_2, \quad p_4 - p_1 = \epsilon_3, \quad p_3 - p_2 = \epsilon_4.$$

Then

$$\epsilon_1 + \epsilon_2 = p_2 - p_1 + p_3 - p_1 = p_2 - p_1 + p_4 - p_2 = p_4 - p_1 = \epsilon_3, \quad (5)$$

since, by Eq. 4, $p_3 - p_1 = p_4 - p_2$.

$$\text{Also,} \quad \epsilon_2 - \epsilon_1 = p_3 - p_2 = \epsilon_4. \quad (6)$$

Hence, if two vectors are drawn outwards from a point, and the parallelogram of which these are two adjacent sides is completed, then the two diagonals of this parallelogram will represent respectively the sum and difference of the two vectors, the sum being that diagonal which passes through the origin of the two vectors, and the difference that which passes through their extremities.*

Again, $p_2 - p_1 + p_3 - p_2 + p_1 - p_3 = 0 = \epsilon_1 + \epsilon_2 + (-\epsilon_3)$; hence the sum of three vectors represented by the sides of a triangle taken around in order is zero.

Similarly, if p_1, p_2, \dots, p_n be any n points whatever taken as corners of a closed polygon, we shall have

$(p_2 - p_1) + (p_3 - p_2) + (p_4 - p_3) + \dots + (p_n - p_{n-1}) + (p_1 - p_n) = 0$; that is, the sum of vectors represented by the sides taken in order about the polygon is zero. By "taken in order" is not meant that any particular order of the points must be observed in forming the polygon, which is evidently unnecessary, but simply that, when the polygon is formed, the vectors will be the operators that will move a point from the starting position along the successive sides back to this position again, so that the final distance from the starting-point will be nothing.

ART. 3. SUM OF TWO WEIGHTED POINTS.†

Consider the sum $m_1 p_1 + m_2 p_2$, in which m_1 and m_2 are scalars, that is, numbers, positive or negative, and p_1, p_2 are points.

* Grassmann (1844), § 15.

† Grassmann (1844), § 95, and (1862), Art. 227.

The scalars m_1 and m_2 will be regarded as values or weights assigned to the points p_1 and p_2 . When any weight is of unit value the figure 1 will be omitted, so that p means $1p$, and is called a unit point. Occasionally, however, a letter may be used to denote a point whose weight is not unity.

To assist his thinking, the reader may consider the weights initially as like or unlike parallel forces acting at the points.

In order to arrive at a meaning for the above expression we shall make two reasonable assumptions, which will prove to be consistent with those already made, viz., first, that the sum is a point, and second, that its weight is the sum of the weights of the two given points. Denoting this sum-point by \bar{p} , we write

$$m_1 p_1 + m_2 p_2 = (m_1 + m_2) \bar{p}. \quad (7)$$

Transposing, we have $m_1(p_1 - \bar{p}) = m_2(\bar{p} - p_2)$, or

$$\frac{p_1 - \bar{p}}{m_2} = \frac{\bar{p} - p_2}{m_1}. \quad (8)$$

Both members of (8) are vectors, and, being equal, they must, by Art. 4, be parallel. This requires that \bar{p} shall be collinear with p_1 and p_2 . Also, since $p_1 - \bar{p}$ and $\bar{p} - p_2$ are vectors whose lengths are respectively the distances from p_1 to \bar{p} and from \bar{p} to p_2 , it follows that these distances are in the ratio of m_2 to m_1 . Hence, \bar{p} is a point on the line $p_1 p_2$ whose distances from p_1 and p_2 are inversely proportional to the weights of these points. We shall call \bar{p} the mean point of the two weighted points. If m_1 and m_2 are both positive, (8) shows that \bar{p} must lie between p_1 and p_2 ; but if one, say m_2 , is negative, let $m_2 = -m_2'$. Thus

$$m_1(p_1 - \bar{p}) = m_2'(p_2 - \bar{p}), \quad (9)$$

and \bar{p} is on the same side of each point, that is, its direction from each point is the same. Also, since its distances from the two points are inversely as their weights, \bar{p} must be nearest the point whose weight is greatest.

Case when $m_1 + m_2 = 0$, or $m_2 = -m_1$.*—With this condition equations (7) and (8) become

$$m_1 p_1 + m_2 p_2 = m_1(p_1 - p_2) = 0 \cdot \bar{p}, \quad (10)$$

and

$$\bar{p} - p_1 = \bar{p} - p_2. \quad (11)$$

Thus \bar{p} is in the same direction from each point, that is, not between them, and yet is equidistant from them. This requires either that the two points shall coincide, that is, $p_2 = p_1$, which evidently satisfies (10) and (11); or else, p_1 and p_2 being different points, that \bar{p} shall be at an infinite distance. Thus the sum is in this case a point of zero weight at infinity.† Eq. (10) shows that a zero point at infinity is equivalent to a vector, or directed quantity, as stated in Art. 1. It has been shown in Art. 2 that $p_2 = p_1$ is the condition that p_1 and p_2 coincide; let us consider the equality of weighted points in general, say $m_1 p_1 = m_2 p_2$. Hence, by (7), there is found $m_1 p_1 - m_2 p_2 = (m_1 - m_2) \bar{p} = 0$; hence, since \bar{p} cannot be zero, $m_1 - m_2 = 0$, or $m_1 = m_2$; and therefore $m_1(p_1 - p_2) = 0$, or, since $m_1 \geq 0$, $p_1 - p_2 = 0$, that is, $p_1 = p_2$. Therefore, if any two points are equal, their weights must be the same and their positions identical, that is, they are the same point.

Exercise 1.—To find the sum and difference of the two weighted points $3p_1$ and p_2 :

$$3p_1 + p_2 = 4\bar{p}, \quad 3p_1 - p_2 = 2\bar{p}',$$

and the mean points are as shown in the figure. The reciprocals of the distances of \bar{p} , p_1 , and \bar{p}' from p_2 ,

viz., $\frac{1}{3}$, $\frac{1}{4}$, $\frac{1}{6}$, are in arithmetical progression, hence the points form a harmonic range.


Exercise 2.—Given a circular disk with a circular disk of

* Grassmann (1862), Art. 222.

† Compare the case of the resultant of unlike parallel forces of equal magnitude.

half its radius removed, as in the figure; to find the centroid of the remaining portion.

Take p_1 at center of large circle, p_3 at center of small circle, and p_2 at the point of contact; then $p_3 = \frac{1}{2}(p_1 + p_2)$. The areas of the two circles are as 1 : 4; call them 1 and 4. Then it is as if there were a weight 4 at p_1 , and a weight -1 at p_3 ; hence $\bar{p} = [4p_1 - \frac{1}{2}(p_1 + p_2)] \div 3 = (7p_1 - p_2) \div 6$.



Prob. 1. Show that $p_1, p_2, m_1p_1 + m_2p_2$, and $m_1p_1 - m_2p_2$ are four points forming a harmonic range.

Prob. 2. An inscribed right-angled triangle is cut from a circular disk; show that the centroid of the remainder of the disk is at the point

$$\frac{(3\pi - 2 \sin 2\alpha) p_1 - p_2 \sin 2\alpha}{3(\pi - \sin 2\alpha)},$$

if p_1 is the center of the circle, p_2 the opposite vertex of the triangle, and α one of its angles.

ART. 4. SUM OF ANY NUMBER OF POINTS.

As in the last article we assume the sum to be a point whose weight is equal to the sum of the weights of the given points; thus,

$$\sum_1^n mp = \bar{p} \sum_1^n m. \quad (12)$$

Let e be some fixed point, and subtract $e \sum_1^n m$ from both sides of (12); thus we have

$$\sum_1^n m(p - e) = (\bar{p} - e) \sum_1^n m, \quad (13)$$

an equation which gives a simple construction for \bar{p} .

If $\sum_1^n m = 0$, then $m_1 = -\sum_2^n m$, and

$$\sum_1^n mp = m_1 p_1 + \sum_2^n mp = m_1 \left(p_1 - \frac{\sum_2^n mp}{\sum_2^n m} \right), \quad (14)$$

so that the sum becomes the difference of two unit points, or a vector whose direction is parallel to the line joining p_1 with the mean of all the other points of the system, and whose length is m_1 times the distance between these points. Since any point of the system may be designated as p_1 , it follows that the line joining any point of the system to the mean of all the others is parallel to any other such line. If $\sum_1^n m p = 0$, equation (14) shows that p is the mean of all the other points of the system, and, since any one of the points may be taken as p_1 , any point of the system is the mean of all the others.

Let $n = 3$ in (12) and (13); then

$$m_1 p_1 + m_2 p_2 + m_3 p_3 = (m_1 + m_2 + m_3) \bar{p}, \quad (15)$$

$$m_1(p_1 - e) + m_2(p_2 - e) + m_3(p_3 - e) = (m_1 + m_2 + m_3)(\bar{p} - e), \quad (16)$$

and \bar{p} is on the line joining the point $m_1 p_1 + m_2 p_2$ with p_3 , and therefore inside the triangle $p_1 p_2 p_3$ if the m 's are all positive. If m_3 be negative and numerically less than $m_1 + m_2$, then \bar{p} will have passed across the line $p_1 p_2$ to the outside of the triangle. If m_1 and m_2 are negative and their sum numerically less than m_3 , then \bar{p} will have passed outside the triangle through p_3 , i.e., it will have crossed $p_2 p_3$ and $p_3 p_1$. The point e must evidently always be in the plane $p_1 p_2 p_3$.

As a numerical example let $m_1 = 3$, $m_2 = 4$, $m_3 = -5$, so that (16) becomes


$$\bar{p} - e = \frac{3}{2}(p_1 - e) + 2(p_2 - e) - \frac{5}{2}(p_3 - e).$$

Now, since e may be any point whatever, put $e = p_3$; then $\bar{p} - p_3 = \frac{3}{2}(p_1 - p_3) + 2(p_2 - p_3)$, and the construction is shown in the figure. $p_4 - p_3 = \frac{3}{2}(p_1 - p_3)$, and $\bar{p} - p_4 = 2(p_2 - p_3)$.

As another example take $\bar{p} = 4p_1 + 5p_2 - 2p_3 - 6p_4$, or, by (13), making $e = p_4$,

$$\begin{aligned} \bar{p} - p_4 &= 4(p_1 - p_4) + 5(p_2 - p_4) - 2(p_3 - p_4) \\ &= p_5 - p_4 + p_6 - p_5 + \bar{p} - p_6. \end{aligned}$$

When any number of geometric quantities can be connected with each other by an equation of the form $\sum m p = 0$, in which the m 's are finite and different from zero, then they are said to be mutually dependent, that is, any one can be expressed in terms of the others. If no such relation can exist between the



quantities, they are independent. We obtain from what has preceded the following conditions:

That two points shall coincide,

$$m_1 p_1 + m_2 p_2 = 0. \quad (17)$$

That three points shall be collinear,

$$m_1 p_1 + m_2 p_2 + m_3 p_3 = 0. \quad (18)$$

That four points shall be coplanar,

$$m_1 p_1 + m_2 p_2 + m_3 p_3 + m_4 p_4 = 0. \quad (19)$$

It follows that three non-collinear points cannot be connected by an equation like (18) unless each coefficient is separately zero. Similarly four non-coplanar points cannot be connected by an equation like (19) unless each coefficient is separately zero.

The significance of these statements will be presently illustrated.

The following are corresponding equations of condition for vectors:

That two vectors shall be parallel,

$$n_1 e_1 + n_2 e_2 = 0. \quad (20)$$

relation of the form of equation (18); hence, as was there shown, each coefficient must be zero; accordingly

$$x_0 - y_1 n_0 = x_1 - y_0 n_1 = y_0 + y_1 = 0,$$

whence we find $x_0 : x_1 = n_0 : -n_1$.

hence $(n_0 - n_1)p_2 = n_0 e_0 - n_1 e_1$, and similarly

$$(n_1 - n_2)p_0 = n_1 e_2 - n_2 e_2, \quad (n_2 - n_0)p_1 = n_2 e_2 - n_0 e_0.$$

Adding, we have

$$(n_1 - n_2)p_0 + (n_2 - n_0)p_1 + (n_0 - n_1)p_2 = 0;$$

therefore, by (18), p_0, p_1, p_2 are collinear.

Exercise 4.—Let $p = \sum_0^2 n e \div \sum_0^2 n$ be any point in the plane of the triangle $e_0 e_1 e_2$: show that lines through the middle points of the sides $e_1 e_2, e_2 e_0$, and $e_0 e_1$ of the triangle parallel to $e_0 p, e_1 p$, and $e_2 p$ meet in a point

$$p' = [(n_1 + n_2)e_0 + (n_2 + n_0)e_1 + (n_0 + n_1)e_2] \div 2 \sum_0^2 n.$$

By the conditions the vector from the middle point of $e_1 e_2$ to p' is a multiple of the vector $e_0 - p$; hence

$$p' - \frac{1}{2}(e_1 + e_2) = x(e_0 - p) \quad \text{or}$$

$$p' = \frac{1}{2}(e_1 + e_2) + x(e_0 - p) = \frac{1}{2}(e_0 + e_1) + y(e_2 - p),$$

or, substituting value of p ,

$$p' = \frac{1}{2}(e_1 + e_2) + x(e_0 - \sum n e \div \sum n) = \frac{1}{2}(e_0 + e_1) + y(e_2 - \sum n e \div \sum n).$$


$$\begin{aligned} \text{hence} \quad & [(x - \frac{1}{2})\sum n + n_0(y - x)]e_0 + n_1(y - x)e_1 \\ & + [(\frac{1}{2} - y)\sum n + n_2(y - x)]e_2 = 0; \end{aligned}$$

therefore, as in the previous exercise, each coefficient must be zero, whence $x = y = \frac{1}{2}$, and substituting we find p' as above. It follows also that the distances of p' from the middle points of the sides are the halves of the distances of p from the opposite vertices.

Prob. 3. Show that $\bar{e} = \frac{1}{3} \sum_0^2 e$ is collinear with p and p' of Exer-


cise 4. Also that, by properly choosing p , it follows that \bar{c} is collinear with the common point of the perpendiculars from the vertices on the opposite sides, and the common point of the perpendiculars to the sides at their middle points.

Prob. 4. Given two circles and an ellipse, as in the figure, with centers at e_0, p_2 , and p_1 . Radii of circles 4 and 1, axes of ellipse 2 and 4, small circle and ellipse touching large circle at e_2 and e_1 , respectively, $e_0e_1e_2$ an equilateral triangle: show that the centroid of the remainder of the large circle, after the small areas are removed, will be at



$$\bar{p} = \frac{1}{3}(16e_0 - p_2 - 2p_1) = \frac{1}{3}(59e_0 - 4e_1 - 3e_2).$$

Prob. 5. If $\frac{4}{9}$ of a sheet of tin in the shape of an isosceles triangle be folded over as in the figure, show that its centroid is given by



$$3\bar{p} = \frac{1}{2}[35(e_0 + e_1) + 11e_2].$$

Prob. 6. If a tetrahedron $e_0e_1e_2e_3$ have a tetrahedron of $\frac{1}{8}$ of its volume cut off by a plane parallel to $e_0e_1e_2$, and one of $\frac{1}{64}$ of its volume cut off by a plane parallel to $e_1e_2e_3$, show that the centroid of the remaining solid is at

$$\bar{p} = \frac{1}{880}(227e_0 + 175e_3 + 239(e_1 + e_2)).$$

ART. 5. REFERENCE SYSTEMS.

Let p be any unit point, e_0, e_1, e_2 three fixed unit points, and w, x, y scalars; then, writing

$$p = we_0 + xe_1 + ye_2, \quad (24)$$

we must have also, because p is a unit point,

$$w + x + y = 1, \quad (25)$$

and p is the mean of the weighted points we_0, xe_1, ye_2 . The point p may occupy any position whatever in the plane $e_0e_1e_2$; for it is on the line joining $we_0 + xe_1$ with e_2 , and by varying y and $w + x, \frac{w}{x}$ remaining constant, p may be moved along

this line from $-\infty$ to $+\infty$; while by varying the ratio $\frac{w}{x}$ the point $w e_0 + x e_1$ may be moved from $-\infty$ to $+\infty$ along $e_0 e_1$, and thus the first line will be rotated through 180 degrees, and p may thus be given any position whatever in the plane.

A system of unit points to which the positions of other points may be referred is called a reference system, and the triangle $e_0 e_1 e_2$ is a reference triangle. For reasons that will appear later, the double area of this triangle will be taken as the unit of measurement of area for a point system in two-dimensional space.

Similarly, in solid space, taking a fourth point e_3 , we write

$$p = w e_0 + x e_1 + y e_2 + z e_3, \quad (26)$$

$$\text{which implies also } w + x + y + z = 1; \quad (27)$$

and p may be shown as above to be capable of occupying any position whatever in space by properly assigning the values of w, x, y, z ; so that e_0, \dots, e_3 form a reference system for points in three-dimensional space. The tetrahedron $e_0 e_1 e_2 e_3$ is called the reference tetrahedron, and six times its volume will be taken as the unit of volume for a point system in three-dimensional space.

Eliminating w between (24) and (25), we have

$$p = e_0 + x(e_1 - e_0) + y(e_2 - e_0), \quad (28)$$

from which it may also be easily seen that p may be any point in the plane $e_0 e_1 e_2$. Writing $p - e_0 = \rho$, $e_1 - e_0 = \epsilon_1$, $e_2 - e_0 = \epsilon_2$,

$$(28) \text{ becomes } \rho = x \epsilon_1 + y \epsilon_2, \quad (29)$$

and ϵ_1, ϵ_2 form a plane reference system for vectors.

Similarly, from (26) and (27) we find

$$\rho = x \epsilon_1 + y \epsilon_2 + z \epsilon_3, \quad (30)$$

and $\epsilon_1, \epsilon_2, \epsilon_3$ are a reference system for vectors in solid space, any vector whatever being expressible in terms of these three.

If, in equations (29) and (30), the reference vectors are of

unit length and mutually perpendicular, we have unit, normal reference systems, and in this case $\iota_1, \iota_2, \iota_3$ will generally be used instead of $\epsilon_1, \epsilon_2, \epsilon_3$.

Exercise 5.—To change from one reference system to another, say from c_0, c_1, c_2 to c'_0, c'_1, c'_2 .

The new reference points must be connected with the old ones by equations such as

$$\begin{aligned} c_0 &= l_0 c'_0 + l_1 c'_1 + l_2 c'_2, & c_1 &= m_0 c'_0 + m_1 c'_1 + m_2 c'_2, \\ c_2 &= n_0 c'_0 + n_1 c'_1 + n_2 c'_2. \end{aligned}$$

Then any point $p = x_0 c_0 + x_1 c_1 + x_2 c_2$ will be expressed in terms of the new reference points by substituting the values of c_0 , etc., as given. If c'_0, c'_1, c'_2 are given in terms of the old points, c_0, c_1, c_2 may be found by elimination. Thus, if $c'_0 = \sum l_e$, $c'_1 = \sum m_e$, $c'_2 = \sum n_e$, we have at once

$$\begin{vmatrix} l_0 & l_1 & l_2 \\ m_0 & m_1 & m_2 \\ n_0 & n_1 & n_2 \end{vmatrix} c_0 = \begin{vmatrix} c'_0 & l_1 & l_2 \\ c'_1 & m_1 & m_2 \\ c'_2 & n_1 & n_2 \end{vmatrix},$$

with similar values for c_1 and c_2 .

As a numerical example let the new reference triangle be formed by joining the middle points of the sides of the old one. Then $c'_0 = \frac{1}{2}(c_1 + c_2)$, $c'_1 = \frac{1}{2}(c_2 + c_0)$, $c'_2 = \frac{1}{2}(c_0 + c_1)$; whence $c_0 = -c'_0 + c'_1 + c'_2$, $c_1 = c'_0 - c'_1 + c'_2$, $c_2 = c'_0 + c'_1 - c'_2$. Thus $p = x_0 c_0 + x_1 c_1 + x_2 c_2$

$$= (-x_0 + x_1 + x_2)c'_0 + (x_0 - x_1 + x_2)c'_1 + (x_0 + x_1 - x_2)c'_2.$$

Exercise 6.—Three points being given in terms of the reference points c_0, c_1, c_2 , find the condition that must hold between their weights when they are collinear.

Let $p_0 = \sum_0^2 l_e c_e$, $p_1 = \sum_0^2 m_e c_e$, $p_2 = \sum_0^2 n_e c_e$; then, k_0, k_1, k_2 being scalars, we must have for collinearity, by (18),

$$k_0 p_0 + k_1 p_1 + k_2 p_2 = 0,$$

that is, $k_0 \sum l e + k \sum_1 m e + k \sum n e = 0$,

whence $(k_0 l_0 + k_1 m_0 + k_2 n_0) e_0 + (k_0 l_1 + k_1 m_1 + k_2 n_1) e_1$
 $+ (k_0 l_2 + k_1 m_2 + k_2 n_2) e_2 = 0$,

and, as e_0, e_1, e_2 are not collinear, the coefficients must be zero, by Art. 4; hence

$$k_0 l_0 + k_1 m_0 + k_2 n_0 = k_0 l_1 + k_1 m_1 + k_2 n_1 = k_0 l_2 + k_1 m_2 + k_2 n_2 = 0,$$

and, by elimination of the k 's,

$$\begin{vmatrix} l_0 & m_0 & n_0 \\ l_1 & m_1 & n_1 \\ l_2 & m_2 & n_2 \end{vmatrix} = 0, \quad (31)$$

which is the required condition of collinearity.

Prob. 7. If $p = 3e_0 - e_1 - e_2$, $4e_0' = 3e_1 + e_2$, $4e_1' = 3e_2 + e_0$, $4e_2' = 3e_0 + e_1$, show that $7p = -19e_0' - 3e_1' + 29e_2'$.

Prob. 8. Find the condition that four points $\sum_0^3 k e, \sum_0^3 l e, \sum_0^3 m e, \sum_0^3 n e$ shall be coplanar. Ans. $[k_0, l_1, m_2, n_3] = 0$.

Prob. 9. If $p = w e_0 + x e_1 + y e_2$, and there exist between the scalars w, x, y a linear relation such as $A w + B x + C y = 0$, A, B, C being scalar constants, show that p will always lie on a straight line which cuts the reference lines in $A e_1 - B e_0$, $A e_2 - C e_0$, and $C e_1 - B e_2$. Consider the special cases when $A = B$, $B = C$, $C = A$, $A = B = C$, $A = 0$, $B = 0$, and $C = 0$.

Prob. 10. If $p = w e_0 + x e_1 + y e_2 + z e_3$, and there exist also an equation $A w + B x + C y + D z = 0$, show that p will lie on a plane which cuts the edges of the reference tetrahedron in $\frac{e_1}{B} - \frac{e_0}{A}$, $\frac{e_2}{C} - \frac{e_0}{A}$, etc. Also, if a second relation between the variables, such as $A' w + B' x + C' y + D' z = 0$, be given, then p lies on a line which pierces the faces of the reference tetrahedron in

$$\begin{vmatrix} e_0 & e_1 & e_2 \\ A & B & C \\ A' & B' & C' \end{vmatrix}, \quad \begin{vmatrix} e_3 & e_0 & e_1 \\ D & A & B \\ D' & A' & B' \end{vmatrix}, \quad \text{etc.}$$

ART. 6. NATURE OF GEOMETRIC MULTIPLICATION.*

The fundamental idea of geometric multiplication is, that a product of two or more factors is that which is determined by those factors.

Thus, two points determine a line passing through them, and also a length, viz., the shortest distance between them; hence $p_1 p_2 = L$ is the sect † drawn from p_1 to p_2 , or generated by a point moving rectilinearly from p_1 to p_2 .

The student should note carefully the difference between $p_1 p_2$ and $p_2 - p_1$; they have the same length and direction, but the sect $p_1 p_2$ is confined to the line through these two points, while the vector $p_2 - p_1$ is not. The sect has position in addition to the direction and length possessed by the vector.

Again, in plane space, two sects determine a point, the intersection of the lines in which they lie, and also an area, as will appear later, so that $L_1 L_2 = p$, in which p is not in general a unit point. In solid space, however, two lines do not, in general, meet, and hence cannot fix a point; but two sects, in this case, determine a tetrahedron of which they are opposite edges.

It appears, therefore, that a product may have different interpretations in spaces of different dimensions. Hence we will consider separately products in plane space, or planimetric products, and those in solid space, or stereometric products.

Products of the kind here considered are termed "combinatory," because two or more factors combine to form a new quantity different from any one of them. This is the fundamental difference between this algebra and the linear associative algebras of Peirce, of which quaternions are a special case.

Before discussing in detail the various products that may arise, we will give a table which will serve as a sort of bird's-eye view of the subject.

* Grassmann (1844), Chap. 2; (1862), Chap. 2.

† See Art. 1.

In this table and generally throughout the chapter we shall use p, p_1, p_2 , etc., for points; $\epsilon, \epsilon_1, \epsilon_2$, etc., for vectors; L, L_1 , etc., for sects, or lines; η, η_1 , etc., for plane-vectors; and P, P_1 , etc., for plane-sects, or planes. Also p, p_1 , etc., as used in this table will not generally be unit points.

The products are arranged in two columns, so as to bring out the geometric principle of duality.

PLANIMETRIC PRODUCTS.

$p_1 p_2 = L.$	$L_1 L_2 = p.$
$p_1 p_2 p_3 = \text{area (scalar).}$	$L_1 L_2 L_3 = (\text{area})^2 (\text{scalar}).$
$pL = \text{area (scalar).}$	$Lp = \text{area (scalar).}$
$p_1 \cdot L_1 L_2 = L.$	$L_1 \cdot p_1 p_2 = p.$
$p_1 p_2 \cdot p_3 p_4 = p.$	$L_1 L_2 \cdot L_3 L_4 = L.$
$p_1 p_2 \cdot p_3 p_4 \cdot p_5 p_6 = (\text{area})^2 (\text{scalar}).$	$L_1 L_2 \cdot L_3 L_4 \cdot L_5 L_6 = (\text{area})^4 (\text{scalar}).$
$\epsilon_1 \epsilon_2 = \text{area (scalar).}$	

STEREOMETRIC PRODUCTS.

$p_1 p_2 = L.$	$P_1 P_2 = L.$
$p_1 p_2 p_3 = P.$	$P_1 P_2 P_3 = p.$
$p_1 p_2 p_3 p_4 = \text{volume (scalar).}$	$P_1 P_2 P_3 P_4 = (\text{volume})^3 (\text{scalar}).$
$pP = \text{volume (scalar).}$	$Pp = \text{volume (scalar).}$
$L_1 L_2 = \text{volume (scalar).}$	$L_1 L_2 = \text{volume (scalar).}$
$pL = Lp = P.$	$PL = LP = p.$
$p \cdot P_1 P_2 = P.$	$P \cdot p_1 p_2 = p.$
$p \cdot P_1 P_2 P_3 = L.$	$P \cdot p_1 p_2 p_3 = L.$
$L \cdot p_1 p_2 p_3 = p.$	$L \cdot P_1 P_2 P_3 = P.$
$\epsilon_1 \epsilon_2 = \eta.$	$\eta_1 \eta_2 = \epsilon.$
$\epsilon_1 \epsilon_2 \epsilon_3 = \text{volume (scalar).}$	$\eta_1 \eta_2 \eta_3 = (\text{volume})^2 (\text{scalar}).$
$\epsilon_1 \epsilon_2 \cdot \epsilon_3 \epsilon_4 = \epsilon.$	$\eta_1 \eta_2 \cdot \eta_3 \eta_4 = \eta.$

Laws of Combinatory Multiplication.—All combinatory products are assumed to be subject to the distributive law expressed by the equation

$$A(B + C) = AB + AC.$$

The planimetric product of three points or of three lines, and the stereometric product of three points or planes, or of four points or planes, are subject to the associative law. That is,

In Plane Space :

$$p_1 p_2 p_3 = p_1 p_2 \cdot p_3 = p_1 \cdot p_2 p_3; \quad L_1 L_2 L_3 = L_1 L_2 \cdot L_3 = L_1 \cdot L_2 L_3.$$

In Solid Space :

$$p_1 p_2 p_3 = p_1 \cdot p_2 p_3 = p_1 p_2 \cdot p_3; \quad P_1 P_2 P_3 = P_1 \cdot P_2 P_3 = P_1 P_2 \cdot P_3.$$

$$p_1 p_2 p_3 p_4 = p_1 \cdot p_2 p_3 p_4 = p_1 p_2 \cdot p_3 p_4;$$

$$P_1 P_2 P_3 P_4 = P_1 \cdot P_2 P_3 P_4 = P_1 P_2 \cdot P_3 P_4.$$

The commutative law of scalar algebra does not, in general, hold. Instead of this, in the products just given as being associative, a law prevails which may be expressed by the equation

$$AB = -BA,$$

from which it follows that the interchange of any two single factors of those products changes the sign of the product.*

Since vectors are equivalent to points at ∞ , the associative law holds for $\epsilon_1 \epsilon_2 \epsilon_3$ and $\eta_1 \eta_2 \eta_3$.

ART. 7. PLANIMETRIC PRODUCTS.

Product of Two Points.†—This has been fully defined in Art. 6, and it is evident from its nature as there given that

$$p_1 p_2 = -p_2 p_1. \quad (32)$$

If $p_2 = p_1$, this becomes $p_1 p_1 = 0$, which must evidently be true, since the sect is now of no length.

$$\text{Also,} \quad p_1(p_2 - p_1) = p_1 p_2 - p_1 p_1 = p_1 p_2. \quad (33)$$

* Grassmann (1862), Chap. 3. † Grassmann (1862), Arts. 245, 246, 247.

But $p_2 - p_1$ is a vector, say, ϵ ; hence

$$p_1 \epsilon = p_1 p_2; \quad (34)$$

or the product of a point and a vector is a sect having the direction and magnitude of the vector; or, again, multiplying a vector by a point fixes its position by making it pass through the point.

To find under what conditions pp' will be equal to $p_1 p_2$. Take any other point p_3 in the plane space under consideration, and write $p = x_1 p_1 + x_2 p_2 + x_3 p_3$, $p' = y_1 p_1 + y_2 p_2 + y_3 p_3$, with the conditions for unit points $\sum x = \sum y = 0$.

$$\text{Then } pp' = \begin{vmatrix} x_1 & x_2 \\ y_1 & y_2 \end{vmatrix} p_1 p_2 + \begin{vmatrix} x_2 & x_3 \\ y_2 & y_3 \end{vmatrix} p_2 p_3 + \begin{vmatrix} x_3 & x_1 \\ y_3 & y_1 \end{vmatrix} p_3 p_1.$$

If this is to reduce to $p_1 p_2$, we must have the third condition $x_2 y_3 - x_3 y_2 = x_3 y_1 - x_1 y_3 = 0$, which requires that $x_3 = y_3 = 0$, unless the coefficient of $p_1 p_2$ is to vanish also. Thus pp' must be in the same straight line with $p_1 p_2$. If, moreover, in addition $x_1 y_2 - x_2 y_1 = 1$, we shall have $pp' = p_1 p_2$. Hence pp' is equal to $p_1 p_2$ when, and only when, the four points are collinear, and p' is distant from p by the same amount and in the same direction that p_2 is from p_1 .

Product of Three Points.—By Art. 6 the product is what is determined by the three points. In solid space they would fix a plane, but, as we are now confined to plane space, this is not the case. The points evidently fix either a triangle or a parallelogram of twice its area, and the product $p_1 p_2 p_3$ will be taken as the area of this, or an equivalent, parallelogram.

This area is taken rather than that of the triangle, because it is what is generated by $p_1 p_2$ as it is moved parallel to its initial position till it passes through p_3 .

We have $p_1 p_2 p_3 = p_1 \cdot p_2 p_3 = -p_1 \cdot p_3 p_2 = -p_1 p_3 p_2$, so that if we go around the triangle in the opposite sense the sign is changed. As this product possesses only the properties of magnitude and sign it is scalar.

$$\text{Write } p = \sum_1^3 x p, p' = \sum_1^3 y p, p'' = \sum_1^3 z p; \text{ then}$$

$$pp'p'' = \begin{vmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{vmatrix} p_1 p_2 p_3; \quad (35)$$

that is, any triple point product in plane space differs from any other only by a scalar factor.*

$$\text{Finally, } p_1 p_2 p_3 = p_1(p_2 - p_1)(p_3 - p_1) = p_1 \epsilon \epsilon', \quad (36)$$

if $\epsilon = p_2 - p_1$ and $\epsilon' = p_3 - p_1$.

Product of Two Vectors.—Using the values of ϵ and ϵ' just given, we see that ϵ and ϵ' determine the same parallelogram that p_1 , p_2 , and p_3 do; hence the meaning of the product is the same in all respects in two-dimensional space.

We shall have $\epsilon \epsilon' = -\epsilon' \epsilon$, for

$$\epsilon \epsilon' = (p_2 - p_1)(p_3 - p_1) = -(p_3 - p_1)(p_2 - p_1) = -\epsilon' \epsilon;$$


since we have shown that inverting the order changes the sign in a product of points. The result may be obtained also by regarding ϵ and ϵ' as points at infinity, or by consideration of a figure.

As we have seen that $\epsilon \epsilon'$ has, in plane space, precisely the same meaning as $p_1 p_2 p_3$ we may write

$$\begin{aligned} p_1 p_2 p_3 &= p_1 \epsilon \epsilon' = \epsilon \epsilon' \\ &= (p_2 - p_1)(p_3 - p_1) = p_2 p_3 + p_3 p_1 + p_1 p_2. \end{aligned} \quad (37)$$

Thus the sum of three sects which form the sides of a triangle, all taken in the same sense as looked at from outside the triangle, is equal to the area of the triangle.

Product of Two Sects.—Any two sects in plane space, L_1 , L_2 , determine a point, the intersection of the lines in which they lie, and an area, that of a parallelogram as in the figure. Let p_0 be the intersection,



and take p_1 and p_2 so that $L_1 = p_0 p_1$ and $L_2 = p_0 p_2$. The area

* Grassmann (1862), Art. 255.

determined by L_1 and L_2 is then the same that we have given as the value of $p_0 p_1 p_2$. We write therefore

$$L_1 L_2 = p_0 p_1 \cdot p_0 p_2 = p_0 p_1 p_2 \cdot p_0. \quad (38)$$

The third member of (38) is not to be regarded as derived from the second by ordinary transposition and reassociation of the points, for the associative law does not hold for the four points taken together, since $p_0 p_1 p_0 \cdot p_2 = 0$. The third member simply results from the definition of $L_1 L_2$.^{*} It may be taken as a model form which will be found to apply to several other cases, for instance to (38) when points and lines are interchanged throughout. Thus, if $p_1 = L_0 L_1$ and $p_2 = L_0 L_2$ we have

$$p_1 p_2 = L_0 L_1 \cdot L_0 L_2 = L_0 L_1 L_2 \cdot L_0. \quad (39)$$

For take p_1' and p_2' so that $p_1 p_1' = L_1$ and $p_2 p_2' = L_2$; $p_1 p_2$ is evidently some multiple of L_0 , say nL_0 ; hence

$$\begin{aligned} p_1 p_2 &= nL_0 = \frac{1}{n^2} (p_1 p_2 \cdot p_1 p_1') \cdot (p_1 p_2 \cdot p_2 p_2') \\ &= \frac{1}{n^2} (p_1 p_2 p_1' \cdot p_1) \cdot (p_1 p_2 p_2' \cdot p_2), \text{ by (38),} \\ &= \frac{1}{n^2} \cdot p_1 p_2 p_1' \cdot p_1 p_2 p_2' \cdot p_1 p_2, \text{ because } p_1 p_2 p_1' \text{ and} \\ &\quad p_1 p_2 p_2' \text{ are scalar,} \\ &= \frac{1}{n} \cdot (p_1 p_2 \cdot p_1 p_1' \cdot p_2 p_2') \cdot L_0, \text{ by (38),} \\ &= L_0 L_1 L_2 \cdot L_0, \text{ which was to be proved.} \end{aligned}$$

Product of Three Sects.—The method has just been indicated, but we may also proceed thus: Let the lines be L_0, L_1, L_2 , and let p_0, p_1, p_2 be their common points. Take scalars n_0, n_1, n_2 so that $L_0 = n_0 p_1 p_2$, etc., then

$$\begin{aligned} L_0 L_1 L_2 &= n_0 n_1 n_2 \cdot p_1 p_2 \cdot p_2 p_0 \cdot p_0 p_1 = -n_0 n_1 n_2 \cdot p_2 p_1 \cdot p_2 p_0 \cdot p_0 p_1 \\ &= -n_0 n_1 n_2 \cdot p_2 p_1 p_0 \cdot p_2 p_0 p_1 = n_0 n_1 n_2 (p_0 p_1 p_2)^2. \end{aligned} \quad (40)$$

^{*} Grassmann applies the terms "eingewandt" and "regressiv" to a product of this kind, the first term being used in the *Ausdehnungslehre* of 1844, and the second in that of 1862. See Chapter 3 of the first, and Chapter 3, Art. 94, of the second.

Product of a Point and Two Sects.—Let p be any point and let L_1 and L_2 be as in (38); then

$$pL_1L_2 = p \cdot p_0p_1 \cdot p_0p_2 = p \cdot p_0p_1p_2 \cdot p_0 = p_0p_1p_2 \cdot pp_0. \quad (41)$$

It has been here assumed that $pL_1L_2 = p \cdot L_1L_2$. The product is not associative, for $pL_1 \cdot L_2$ is the line L_2 times the scalar pL_1 , a different meaning from that assigned in (41). As a rule, to avoid ambiguity, the grouping of such products will be indicated by dots.

Product of Two Parallel Sects.—Let them be $p_1\epsilon$ and $n p_2\epsilon$; then, as in (38),

$$p_1\epsilon \cdot n p_2\epsilon = n \cdot p_1\epsilon \cdot p_2\epsilon = n \cdot \epsilon p_1 \cdot \epsilon p_2 = n \cdot \epsilon p_1p_2 \cdot \epsilon, \quad (42)$$

that is, a scalar times the common point at ∞ .

Addition and Subtraction of Sects.—Let L_1 and L_2 be two sects, p_0 their common point, and p_1 and p_2 so taken that $L_1 = p_0p_1$, $L_2 = p_0p_2$; then

$$L_1 + L_2 = p_0p_1 + p_0p_2 = p_0(p_1 + p_2) = 2p_0\bar{p}, \quad (43)$$

\bar{p} being the mean of p_1 and p_2 ; hence the sum is that diagonal of the parallelogram which passes through p_0 . Also

$$L_1 - L_2 = p_0(p_1 - p_2), \quad (44)$$

so that the difference of the two passes also through p_0 and is parallel to the other diagonal of the parallelogram determined by L_1 and L_2 .

If the two sects are parallel let them be $n_1p_1\epsilon$ and $n_2p_2\epsilon$; then

$$n_1p_1\epsilon + n_2p_2\epsilon = (n_1p_1 + n_2p_2)\epsilon = (n_1 + n_2)\bar{p}\epsilon, \quad (45)$$

so that the sum is a sect parallel to each of them, having a length equal to the sum of their lengths, and at distances from them inversely proportional to their lengths.

If $n_2 = -n_1$ the two sects are oppositely directed and of equal length, and the sum is

$$n_1(p_1\epsilon - p_2\epsilon) = n_1(p_1 - p_2)\epsilon, \quad (46)$$

which, being the product of two vectors, is a scalar area.

Multiply the equations by p_1 , p_2 , and p_3 respectively, and add; hence

$$x \sum_1^3 ap + y \sum_1^3 bp + z \sum_1^3 cp = \sum_1^3 dp.$$


Now $\sum ap$, $\sum bp$, etc., are points: multiply the equation just written by $\sum ap$, $\sum bp$; thus

$$z \sum ap \cdot \sum bp \sum cp = \sum ap \cdot \sum bp \cdot \sum dp,$$

because $\sum ap \cdot \sum ap = 0$, etc.; therefore

$$z = \sum ap \cdot \sum bp \cdot \sum dp \div \sum ap \cdot \sum bp \sum cp = [a_1, b_2, d_3] \div [a_1, b_2, c_3],$$

a very simple proof of the determinant solution. Of course x and y will be found by multiplying by the other pairs of points.



Exercise 8.—Forces are represented by given multiples of the sides of a parallelogram; determine their resultant.

Let the parallelogram be double the triangle $e_0e_1e_2$, and the forces

$$\begin{aligned} k_0e_0e_1 + k_1e_1(e_2 - e_0) + k_2e_2(e_0 - e_1) + k_3e_2e_0 &= \sum pe \\ &= (k_0 + k_1)e_0e_1 + (k_1 + k_2)e_1e_2 + (k_2 + k_3)e_2e_0. \end{aligned}$$

Multiply by e_0e_1 to find where the resultant cuts this line, then

$$(k_1 + k_2)e_0e_1 \cdot e_1e_2 + (k_2 + k_3)e_0e_1 \cdot e_2e_0 = e_0e_1e_2 \cdot [(k_1 + k_2)e_1 - (k_2 + k_3)e_0],$$

or e_0e_1 cuts the resultant at the point

$$[(k_1 + k_2)e_1 - (k_2 + k_3)e_0] \div (k_1 - k_3).$$

Similarly the resultant cuts the other sides of the reference triangle at $[(k_2 + k_3)e_2 - (k_0 + k_1)e_1] \div (k_2 + k_3 - k_0 - k_1)$ and at $[(k_0 + k_1)e_0 - (k_1 + k_2)e_2] \div (k_0 - k_2)$.

Suppose $k_0 = k_1 = k_2 = k_3$; then each of the three points just found recedes to infinity; but in this case $\sum pe$ reduces to $2k_0(e_0e_1 + e_1e_2 + e_2e_0) = 2k_0(e_1 - e_0)(e_2 - e_0)$, and the system is equivalent to a couple.

Prob. 11. Construct the resultant of Exercise 8 when $k_0 = 1$, $k_1 = 2$, $k_2 = 3$, $k_3 = 4$; when $k_0 = 1$, $k_1 = -2$, $k_2 = 3$, $k_3 = -4$; when $k_0 = 3$, $k_1 = k_3 = 2$, $k_2 = 1$; and when $k_1 = k_2 = 1$, $k_0 = k_3 = -2$.

Prob. 12. There are given n points p_1, \dots, p_n ; to find a point e such that forces represented by the sects ep_1, ep_2 , etc., shall be in equilibrium. (The equation of equilibrium is $\sum ep \equiv e \sum p \equiv \frac{1}{n} e \bar{p} = 0$. Hence e coincides with the mean point of the p 's.)

Prob. 13. If a harmonic range e_1, p, e_2, p' be given, together with some point e_0 not collinear with these points, show that

$$e_0 e_1 p \cdot e_0 e_2 p' = -e_0 p e_2 \cdot e_0 p' e_1.$$

(Let $p = m_1 e_1 + m_2 e_2$ and $p' = m_1 e_1 - m_2 e_2$, as in Exercise 2 of Art. 3.)

Prob. 14. Show that the relation of Prob. 13 holds for any four points whatever taken respectively on the four lines $e_0 e_1, e_0 p, e_0 e_2, e_0 p'$. If the four points are all at the same distance from e_0 , show that the areas $e_0 e_1 p$, etc., become proportional to the sines of the angles between $e_0 e_1$ and $e_0 p$, etc.

ART. 8. THE COMPLEMENT.*

Taking point reference systems, or unit normal vector reference systems, as in Art. 5, the product of the reference units taken in order being in any case unity, the complement of any reference unit is the product of all the others so taken that the unit times its complement is unity.

To find the complements of quantities other than reference units the following properties are assumed:

(a) The complement of a product is equal to the product of the complements of its factors.

(b) The complement of a sum is equal to the sum of the complements of the terms added together.

(c) The complement of a scalar quantity is the scalar itself.

Considering now the point system in plane space e_0, e_1, e_2 with the constant condition $e_0 e_1 e_2 = 1$, the sides of the reference triangle taken in order are the complements of the opposite vertices, and vice versa.

The complement of a quantity is indicated by a vertical line, as $|p$, read, complement of p .

* See Ausdehnungslehre of 1862, Art. 89.

$$\begin{aligned} \text{Thus} \quad |e_0 = e_1 e_2, \quad |e_1 e_2 = |(|e_0) = e_0, \\ |e_1 = e_2 e_0, \quad |e_2 e_0 = |(|e_1) = e_1, \\ |e_2 = e_0 e_1, \quad |e_0 e_1 = |(|e_2) = e_2. \end{aligned}$$

For $e_0 | e_0 = e_0 e_1 e_2 = 1$, which agrees with the definition ;

$|e_1 e_2 = |e_1 \cdot |e_2 = e_2 e_0 \cdot e_0 e_1 = -e_0 e_2 \cdot e_0 e_1 = -e_0 e_2 e_1 \cdot e_0 = e_0$, by (a) and (38) ;

$|e_0 e_1 e_2 = |e_0 \cdot |e_1 \cdot |e_2 = e_1 e_2 \cdot e_2 e_0 \cdot e_0 e_1 = (e_0 e_1 e_2)^2 = 1 = e_0 e_1 e_2$, which agrees with (c) ; $e_0 | e_1 = e_0 e_2 e_0 = 0 = e_0 | e_2 = e_1 | e_2$.

Next take any point $p_1 = \sum_0^2 l e$, and we have, by (b),

$$|p_1 = \sum_0^2 l |e = l_0 e_1 e_2 + l_1 e_2 e_0 + l_2 e_0 e_1 = l_0 l_1 l_2 \left(\frac{e_1}{l_1} - \frac{e_0}{l_0} \right) \left(\frac{e_2}{l_2} - \frac{e_0}{l_0} \right) = L_1. \quad (50)$$

Thus the complement of a point is a line,* which may be easily constructed by the fourth member of (50), which expresses this line as the product of the points in which it cuts the sides $e_0 e_1$ and $e_0 e_2$ of the reference triangle. Comparing this equation with Ex. 3 in Art. 4, it appears that $|p_1$ above is related to the point $\sum_0^2 \frac{e}{l}$ as the line $p_0 p_2$ of Ex. 3 is to the point $\sum me$. Hence $|p_1$ may be found by constructing this line corresponding to $\sum_0^2 \frac{e}{l}$ as shown in the figure of Ex. 3, Art. 4.

Again, the line $|p_1$ may be shown to be the anti-polar of p with respect to an ellipse of such dimensions, and so placed upon $e_0 e_1 e_2$ that, with reference to it, each side of the reference triangle is the anti-polar of the opposite vertex.* From this it appears that complementary relations are polar reciprocal relations. Take any point $p_2 = \sum_0^2 m e$, and we have

$$\begin{aligned} p_1 | p_2 &= (l_0 e_0 + l_1 e_1 + l_2 e_2) (m_0 e_1 e_2 + m_1 e_2 e_0 + m_2 e_0 e_1) \\ &= \sum_0^2 l m = \sum me \cdot \sum l |e = p_2 | p_1, \end{aligned} \quad (51)$$

* See Hyde's Directional Calculus, Arts 41-43 and 121-123.

so that this product is commutative about the complement sign, and scalar. This is true of all such products when the quantities on each side of the complement sign are of the same order in the reference units. Take for instance the product $p_1 p_2 | p_3 p_4$. This is scalar because $| p_3 p_4$ is a point, so that the whole quantity is equivalent to a triple-point product; and we have $p_1 p_2 | p_3 p_4 = | p_3 p_4 \cdot p_1 p_2 = | (p_3 p_4 | p_1 p_2) = p_3 p_4 | p_1 p_2$, by (a) and (c). If, however, such a quantity be taken as $p_1 p_2 \cdot | p_3$ it is neither scalar nor commutative about the sign $|$; for, $| p_3$ being a line, the product is that of two lines, that is, a point, and

$$p_1 p_2 \cdot | p_3 = - | p_3 \cdot p_1 p_2 = - | (p_3 \cdot | p_1 p_2). \quad (52)$$

Such products as we have just been considering are called by Grassmann "inner products,"* and he regards the sign $|$ as a multiplication sign for this sort of product. Inasmuch, however, as these products do not differ in nature from those heretofore considered, it appears to the author to conduce to simplicity not to introduce a nomenclature which implies a new species of multiplication. For instance, $p | q$ will be treated as the combinatory product of p into the complement of q , and not as a different kind of product of p into q .

The term co-product may be applied to such expressions, regarded as an abbreviation merely, after the analogy of cosine for complement of the sine.

Consider next a unit normal vector system. By the definition we have

$$| l_1 = l_2, \quad | l_2 = | (| l_1) = - l_1,$$


$$\text{because } l_1 | l_1 = l_1 l_2 = I,$$

$$l_2 | l_2 = l_2 (- l_1) = - l_2 l_1 = l_1 l_2 = I.$$

$$\text{Also, } l_1 | l_2 = - l_1 l_1 = 0 = l_2 | l_1.$$

Next let

$$\epsilon_1 = m_1 l_1 + m_2 l_2 \quad \text{and} \quad \epsilon_2 = n_1 l_1 + n_2 l_2;$$



* Grassmann (1862), Chapter 4.

then, by (b) and (c),

$$|\epsilon_1 = m_1|l_1 + m_2|l_2 = m_1l_2 - m_2l_1. \quad (53)$$

By the figure it is evident that $|\epsilon_1$ is a vector of the same length as ϵ_1 and perpendicular to it, or, in other words, taking the complement of a vector in plane space rotates it positively through 90° .

The co-product $\epsilon_1|\epsilon_2$ is the area of the parallelogram, two of whose sides are ϵ_1 and $|\epsilon_2$ drawn outwards from a point; if ϵ_1 is parallel to $|\epsilon_2$, this area vanishes, or $\epsilon_1|\epsilon_2 = 0$; but, since $|\epsilon_2$ is perpendicular to ϵ_2 , ϵ_1 must in this case be perpendicular to ϵ_2 ; hence the equation

$$\epsilon_1|\epsilon_2 = 0 \quad (54)$$

is the condition that two vectors ϵ_1 and ϵ_2 shall be perpendicular to each other.

The co-product $\epsilon_1|\epsilon_1$, which will usually be written ϵ_1^2 , and called the co-square of ϵ_1 , is the area of a square each of whose sides has the length $T\epsilon_1$; hence

$$T\epsilon_1 = \sqrt{\epsilon_1|\epsilon_1} = \sqrt{\epsilon_1^2}. \quad (55)$$

Let α_1 and α_2 be the angles between l_1 and ϵ_1 and between l_1 and ϵ_2 respectively, as in the figure. Then

$$\epsilon_1\epsilon_2 = m_1n_2 - m_2n_1 = T\epsilon_1T\epsilon_2 \sin(\alpha_2 - \alpha_1), \quad (56)$$

the third member being the ordinary expression for the area of the parallelogram $\epsilon_1\epsilon_2$. Also

$$\begin{aligned} \epsilon_1|\epsilon_2 &= (m_1l_1 + m_2l_2)(n_1l_2 - n_2l_1) \\ &= m_1n_1 + m_2n_2 = T\epsilon_1T\epsilon_2 \cos(\alpha_2 - \alpha_1), \end{aligned} \quad (57)$$

the last member being found as before, remembering that $\sin(90^\circ + \alpha_2 - \alpha_1) = \cos(\alpha_2 - \alpha_1)$.

If in (57) we let $\epsilon_2 = \epsilon_1$, whence $n_1 = m_1$ and $n_2 = m_2$, we have

$$T\epsilon_1 = \sqrt{\epsilon_1^2} = \sqrt{m_1^2 + m_2^2}. \quad (58)$$

If $T\epsilon_1 = T\epsilon_2 = 1$, then $m_1 = \cos \alpha_1$, $m_2 = \sin \alpha_1$, $n_1 = \cos \alpha_2$, $n_2 = \sin \alpha_2$, and equations (56) and (57) give the ordinary trigonometrical formulas $\sin(\alpha_2 - \alpha_1) = \sin \alpha_2 \cos \alpha_1 - \cos \alpha_2 \sin \alpha_1$,

and $\cos(\alpha_2 - \alpha_1) = \cos \alpha_1 \cos \alpha_2 + \sin \alpha_1 \sin \alpha_2$. Squaring and adding (56) and (57), there results

$$T^2 \epsilon_1 \cdot T^2 \epsilon_2 = \epsilon_1^2 \epsilon_2^2 = (\epsilon_1 \epsilon_2)^2 + (\epsilon_1 | \epsilon_2)^2. \quad (59)$$

Attention is called to the fact, which the student may have already noticed, that such an equation as $AB = AC$, in which AB and AC are combinatory products, does not, in general, imply that $B = C$, for the reason that the equation $A(B - C) = 0$ can usually be satisfied without either factor being itself zero. Thus $pL_1 = pL_2$ means simply that the two quantities which are equated have the same magnitude and sign, which permits L_2 to have an infinity of lengths and positions, when p and L_1 are given. The equation $p_1 p_2 = p_1 p_3$, or $p_1(p_2 - p_3) = 0$, p_2 and p_3 being unit points, implies, however, that $p_2 = p_3$, unless p_1 is at ∞ , that is, a vector.

Exercise 9.—A triangle whose sides are of constant length moves so that two of its vertices remain on two fixed lines: find the locus of the other vertex.

Let $e_0 \epsilon_1$ and $e_0 \epsilon_2$ be the two fixed lines, and $pp'p''$ the triangle. Let pe be perpendicular to $p'p''$, $p' - e_0 = x\epsilon_1$ and $p'' - e_0 = y\epsilon_2$; then $p'' - p' = y\epsilon_2 - x\epsilon_1$, $T(y\epsilon_2 - x\epsilon_1) = c = \text{constant}$, by the conditions. Also, $Tp'e = \text{constant} = mc$, say, and $Te p = \text{constant} = nc$, say. Hence

$$e - p' = Tp'e. U(e - p') = mc \cdot \frac{y\epsilon_2 - x\epsilon_1}{T(y\epsilon_2 - x\epsilon_1)} = m(y\epsilon_2 - x\epsilon_1),$$

and similarly $p - e = n(y\epsilon_2 - x\epsilon_1)$. Therefore


$$p - e_0 = \rho = x\epsilon_1 + m(y\epsilon_2 - x\epsilon_1) + n(y\epsilon_2 - x\epsilon_1),$$

an equation which, with the condition $T(y\epsilon_2 - x\epsilon_1) = c$, or

$$y^2 \epsilon_2^2 - 2xy\epsilon_1 | \epsilon_2 + x^2 \epsilon_1^2 = c^2,$$

determines the locus to be a second-degree curve, which must in fact be an ellipse, since it can have no points at infinity. Let us rearrange the equation in ρ thus:

$$\rho = x[(1 - m)\epsilon_1 - n|\epsilon_1] + y[m\epsilon_2 + n|\epsilon_2] = x\epsilon + y\epsilon', \text{ say,}$$



so that $\epsilon = (1 - m)\epsilon_1 - n|\epsilon_1$ and $\epsilon' = m\epsilon_2 + n|\epsilon_2$; then multiply successively into ϵ and ϵ' ; therefore $\rho\epsilon = j\epsilon'\epsilon$ and $\rho\epsilon' = x\epsilon\epsilon'$. Substituting these values of x and j in the equation of condition, we have

$$\epsilon_2^2 \cdot (\rho\epsilon)^2 + 2\epsilon_1|\epsilon_2 \cdot \rho\epsilon \cdot \rho\epsilon' + \epsilon_1^2(\rho\epsilon')^2 = \epsilon^2(\epsilon\epsilon')^2,$$


a scalar equation of the second degree in ρ .

Exercise 10.—There is given an irregular polygon of n sides: show that if forces act at the middle points of these sides, proportional to them in magnitude, and directed all outward or else all inward, these forces will be in equilibrium.

Let ϵ_0 be a vertex of the polygon, and let $2\epsilon_1, 2\epsilon_2, \dots, 2\epsilon_n$ represent its sides in magnitude and direction. Then the middle points will be $\epsilon_0 + \epsilon_1, \epsilon_0 + 2\epsilon_1 + \epsilon_2$, etc., and, using the complement in a vector system, we have

$$\begin{aligned} \sum p\epsilon &= (\epsilon_0 + \epsilon_1)|\epsilon_1 + (\epsilon_0 + 2\epsilon_1 + \epsilon_2)|\epsilon_2 + (\epsilon_0 + 2\epsilon_1 + 2\epsilon_2 + \epsilon_3)|\epsilon_3 + \dots \\ &\quad + (\epsilon_0 + 2\epsilon_1 + \dots + 2\epsilon_{n-1} + \epsilon_n)|\epsilon_n \\ &= \epsilon_0 \left| \sum_1^n \epsilon + \sum_1^n \epsilon^2 + 2\epsilon_1 \right| \sum_2^n \epsilon + 2\epsilon_2 \left| \sum_3^n \epsilon + \dots + 2\epsilon_{n-1} \right| \epsilon_n \\ &= \epsilon_0 \left| \sum_1^n \epsilon + \left(\sum_1^n \epsilon \right)^2 \right| = 0, \text{ which was to be proved.} \end{aligned}$$

Exercise 11.—A line passes through a fixed point and cuts two fixed lines; at the points of intersection perpendiculars to the fixed lines are erected; find the locus of the intersection of these perpendiculars.



Let the fixed lines be $c_0\epsilon_1$ and $c_0\epsilon_2$, and the fixed point $c_0 + \epsilon_3$; the moving line cuts the fixed lines in p' and p'' , at which points perpendiculars are erected meeting in p .

Let $p - c_0 = \rho$, $p' - c_0 = x\epsilon_1$, $p'' - c_0 = y\epsilon_2$, $T\epsilon_1 = T\epsilon_2 = 1$; then $\rho = x\epsilon_1 + x'|\epsilon_1 = y\epsilon_2 + y'|\epsilon_2$, whence $\rho|\epsilon_1 = x$ and $\rho|\epsilon_2 = y$.

Also, since $e_0 + e_3, p', p''$ are collinear points,

$$(x\epsilon_1 - e_3)(y\epsilon_2 - e_3) = 0 = xy\epsilon_1\epsilon_2 + y\epsilon_2e_3 + x\epsilon_3e_1;$$

or, substituting values of x and y ,

$$\rho|\epsilon_1 \cdot \rho|\epsilon_2 \cdot \epsilon_1\epsilon_2 + \rho|\epsilon_2 \cdot \epsilon_2\epsilon_3 + \rho|c_1 \cdot \epsilon_3e_1 = 0,$$

an equation of the second degree in ρ , and hence representing a conic.

Prob. 15. If a, b, c are the lengths of the sides of a triangle, prove the formula $a^2 = b^2 + c^2 - 2bc \cos A$, by taking vectors ϵ_1, ϵ_2 , and $\epsilon_2 - \epsilon_1$ equal to the respective sides.

Prob. 16. If $e_0\epsilon_1$ and $e_0\epsilon_2$ are two unit lines, show that the vector perpendicular from e_0 on the line $(e_0 + a\epsilon_1)(e_0 + b\epsilon_2)$ is

$\frac{ab\epsilon_1\epsilon_2}{(b\epsilon_2 - a\epsilon_1)^2} \cdot (b\epsilon_2 - a\epsilon_1)$, of which the length is $\frac{ab\epsilon_1\epsilon_2}{T(b\epsilon_2 - a\epsilon_1)}$. From this derive the Cartesian expression for the perpendicular from the origin upon a straight line in oblique coordinates,
 $ab \sin \omega \div (a^2 + b^2 - 2ab \cos \omega)^{1/2}$, ω being angle between the axes.

Prob. 17. If three points, $me_0 + ne_1, me_1 + ne_2, me_2 + ne_0$, be taken on the sides of the reference triangle, then the sides of the complementary triangle, $(me_0 + ne_1)$, etc., will be respectively parallel to the corresponding sides of the triangle formed by the assumed points $(me_1 + ne_2), (me_2 + ne_0)$, etc.

ART. 9. EQUATIONS OF CONDITION, AND FORMULAS.

Several equations of condition are placed here together for convenient reference: some have been already given; others follow from the results of Arts. 7 and 8. When we have

$$\begin{array}{l|l} \left. \begin{array}{l} p_1 p_2 = 0, \\ \text{or } n_1 p_1 + n_2 p_2 = 0, \end{array} \right\} & \left. \begin{array}{l} L_1 L_2 = 0, \\ \text{or } n_1 L_1 + n_2 L_2 = 0, \end{array} \right\} \quad (60) \\ \text{the two points coincide;} & \text{the two lines coincide;} \end{array}$$

$$\begin{array}{l|l} \left. \begin{array}{l} p_1 p_2 p_3 = 0, \\ \text{or } \sum_1^3 n_i p_i = 0, \end{array} \right\} & \left. \begin{array}{l} L_1 L_2 L_3 = 0, \\ \text{or } \sum_1^3 n_i L_i = 0, \end{array} \right\} \quad (61) \\ \text{the three points are collinear;} & \text{the three lines are confluent.} \end{array}$$

$$\epsilon_1 \epsilon_2 = 0, \quad \text{or} \quad n_1 \epsilon_1 + n_2 \epsilon_2 = 0, \quad (62)$$

the two vectors are parallel (points at infinity coincide);

$$\epsilon_1 | \epsilon_2 = 0, \quad (63)$$

the two vectors are perpendicular ;

$$\left. \begin{array}{l} p_1 | p_2 = 0, \\ \text{either point lies on the com-} \\ \text{plementary line of the other.} \end{array} \right\} \begin{array}{l} L_1 | L_2 = 0, \\ \text{either line passes through the} \\ \text{complementary point of the} \\ \text{other.} \end{array} \quad (64)$$

If we write the equation

$$\rho = x_1 \epsilon_1 + x_2 \epsilon_2,$$

$x_1 \epsilon_1$ is the projection of ρ on ϵ_1 parallel to ϵ_2 , and $x_2 \epsilon_2$ is the projection of ρ on ϵ_2 parallel to ϵ_1 . Multiply both sides of the equation into ϵ_2 ; therefore $\rho \epsilon_2 = x_1 \epsilon_1 \epsilon_2$, or $x_1 = \rho \epsilon_2 \div \epsilon_1 \epsilon_2$. Similarly, multiplying into ϵ_1 , we have $\rho \epsilon_1 = x_2 \epsilon_2 \epsilon_1$, or $x_2 = \rho \epsilon_1 \div \epsilon_2 \epsilon_1$, whence

$$\rho = \frac{\epsilon_1 \cdot \rho \epsilon_2}{\epsilon_1 \epsilon_2} + \frac{\epsilon_2 \cdot \rho \epsilon_1}{\epsilon_2 \epsilon_1}. \quad (65)$$

The two terms of the second member of (65) are therefore the projections of ρ on ϵ_1 parallel to ϵ_2 , and on ϵ_2 parallel to ϵ_1 , respectively.*

Let ϵ_1 and ϵ_2 be unit normal vectors, say, i and j ; then (65) becomes

$$\rho = i \cdot \rho | i - j \cdot \rho | j = i \cdot \rho | i + j \rho \cdot | j; \quad (66)$$

or, if i_1 and i_2 be used instead of i and j ,

$$\rho = i_1 \cdot \rho | i_1 + i_2 \cdot \rho | i_2. \quad (67)$$

Again, in (65) let $\rho = \epsilon_3$, clear of fractions, and transpose ; therefore

$$\epsilon_1 \epsilon_2 \cdot \epsilon_3 + \epsilon_2 \epsilon_3 \cdot \epsilon_1 + \epsilon_3 \epsilon_1 \cdot \epsilon_2 = 0, \quad (68)$$

a symmetrical relation between any three directions in plane space. Let $T \epsilon_1 = T \epsilon_2 = T \epsilon_3 = 1$, and multiply (68) into $|\epsilon_3$, thus

$$\epsilon_1 \epsilon_2 + \epsilon_2 \epsilon_3 \cdot \epsilon_1 | \epsilon_3 + \epsilon_3 \epsilon_1 \cdot \epsilon_2 | \epsilon_3 = 0, \quad (69)$$

which is equivalent to

$$\sin (\alpha \pm \beta) = \sin \alpha \cos \beta \pm \cos \alpha \sin \beta,$$

the upper or lower sign corresponding to the case when ϵ_3 is

* Grassmann (1844), Chapter 5 (1862), Art. 129. Hyde's Directional Calculus, Arts. 46 and 47.

between ϵ_1 and ϵ_2 , or outside, respectively. Writing in (69) $|\epsilon_2$ instead of ϵ_3 , we have

$$\epsilon_1|\epsilon_2 - \epsilon_2|\epsilon_3 \cdot \epsilon_3|\epsilon_1 + \epsilon_3\epsilon_1 \cdot \epsilon_2\epsilon_3 = 0, \quad (70)$$

which gives the $\cos(\alpha \pm \beta)$. These formulas being for any three directions in plane space, are independent of the magnitude of the angles involved.

There is given below a set of formulas for points and lines, arranged in complementary pairs, and all placed together for convenient reference, the derivation of them following after.

$$\begin{aligned} p &= (p_0 p_1 p_2)^{-1} [p_0 \cdot p p_1 p_2 + p_1 \cdot p p_2 p_0 + p_2 \cdot p p_0 p_1], \\ L &= (L_0 L_1 L_2)^{-1} [L_0 \cdot L L_1 L_2 + L_1 \cdot L L_2 L_0 + L_2 \cdot L L_0 L_1] \end{aligned} \quad (71)$$

$$\begin{aligned} p &= (p_0 p_1 p_2)^{-1} [p_1 p_2 \cdot p | p_0 + p_2 p_0 \cdot p | p_1 + p_0 p_1 \cdot p | p_2], \\ L &= (L_0 L_1 L_2)^{-1} [L_1 L_2 \cdot L | L_0 + L_2 L_0 \cdot L | L_1 + L_0 L_1 \cdot L | L_2] \end{aligned} \quad (72)$$

$$\begin{aligned} p_1 p_2 \cdot p_3 p_4 &= -p_1 \cdot p_2 p_3 p_4 + p_2 \cdot p_3 p_4 p_1 \\ &= p_3 \cdot p_4 p_1 p_2 - p_4 \cdot p_1 p_2 p_3, \\ L_1 L_2 \cdot L_3 L_4 &= -L_1 \cdot L_2 L_3 L_4 + L_2 \cdot L_3 L_4 L_1 \\ &= L_3 \cdot L_4 L_1 L_2 - L_4 \cdot L_1 L_2 L_3 \end{aligned} \quad (73)$$

$$p_1 p_2 \cdot |q_1 = - \left| \begin{array}{cc} p_1 & p_1 | q_1 \\ p_2 & p_2 | q_1 \end{array} \right|, \quad L_1 L_2 | M_1 = - \left| \begin{array}{cc} L_1 & L_1 | M_1 \\ L_2 & L_2 | M_1 \end{array} \right|, \quad (74)$$

$$p_2 | q_1 q_2 = \left| \begin{array}{cc} q_1 & p_2 | q_1 \\ q_2 & p_2 | q_2 \end{array} \right|, \quad L_2 | M_1 M_2 = \left| \begin{array}{cc} M_1 & L_2 | M_1 \\ M_2 & L_2 | M_2 \end{array} \right|, \quad (75)$$

$$p_1 p_2 | q_1 q_2 = \left| \begin{array}{cc} p_1 | q_1 & p_1 | q_2 \\ p_2 | q_1 & p_2 | q_2 \end{array} \right|, \quad L_1 L_2 | M_1 M_2 = \left| \begin{array}{cc} L_1 | M_1 & L_1 | M_2 \\ L_2 | M_1 & L_2 | M_2 \end{array} \right|, \quad (76)$$

$$p_0 p_1 p_2 \cdot q_0 q_1 q_2 = \left| \begin{array}{ccc} p_0 | q_0 & p_0 | q_1 & p_0 | q_2 \\ p_1 | q_0 & p_1 | q_1 & p_1 | q_2 \\ p_2 | q_0 & p_2 | q_1 & p_2 | q_2 \end{array} \right| \quad (77)$$

The complementary formula to (77) is not given, but may be obtained by putting L 's and M 's for p 's and q 's.

Derivation of Equations (71)-(77).—Equation (71). Write $p = x_0 p_0 + x_1 p_1 + x_2 p_2$, and multiply this equation by $p_1 p_2$; then $p_1 p_2 p = x_0 p_1 p_2 p_0$, or $x_0 = p p_1 p_2 \div p_0 p_1 p_2$.

Multiplying similarly by $p_0 p_2$ and by $p_0 p_1$, we find $x_1 = p p_2 p_0 \div p_0 p_1 p$ and $x_2 = p p_0 p_1 \div p_0 p p_2$. The substitu-

tion of these values gives the first of (71), and the second is similarly obtained or may be found by simply putting L 's for p 's in the first.

Equation (72). Write $p = x_0 | p_1 p_2 + x_1 | p_2 p_0 + x_2 | p_0 p_1$, and multiply into $| p_0$; thus $p | p_0 = x_0 p_0 p_1 p_2$. Find in the same way values of x_1 and x_2 , and substitute.

Equation (73). Write $p_1 p_2 \cdot p_3 p_4 = x p_1 + y p_2$, and multiply by $p p_2$; therefore $p p_2 \cdot p_1 p_2 \cdot p_3 p_4 = x p p_2 p_1$, or, by Eq. (38), $p_2 p p_1 \cdot p_2 p_3 p_4 = x p p_2 p_1 = -x p_2 p p_1$; or, $x = -p_2 p_3 p_4$. Multiplying by $p p_1$ we find $y = p_3 p_4 p_1$, and on substituting obtain the first of (73). For the second put $p_1 p_2 \cdot p_3 p_4 = x p_3 + y p_4$, and proceed in a similar way.

Equation (74). In the first of (73) put $p_3 p_4 = | q_1$.

Equation (75). In the fourth of (73) put

$$L_1 L_2 = p_2, L_3 = | q_1, L_4 = | q_2.$$

Equation (76). Multiply (75) by p_1 .

Equation (77). In the first of (72) put q_2 for p , and multiply by $p_0 p_1 p_2 \cdot q_0 q_1$; then

$$\begin{aligned} p_0 p_1 p_2 \cdot q_0 q_1 q_2 &= q_0 q_1 | p_1 p_2 \cdot q_2 | p_0 + q_0 q_1 | p_2 p_0 \cdot q_2 | p_1 + q_0 q_1 | p_0 p_1 \cdot q_2 | p_2 \\ &= p_0 | q_2 \cdot \begin{vmatrix} p_1 q_0 & p_1 q_1 \\ p_2 q_0 & p_2 q_1 \end{vmatrix} + p_1 | q_2 \cdot \begin{vmatrix} p_2 q_0 & p_2 q_1 \\ p_0 q_0 & p_0 q_1 \end{vmatrix} + p_2 | q_2 \cdot \begin{vmatrix} p_0 q_0 & p_0 q_1 \\ p_1 q_0 & p_1 q_1 \end{vmatrix}, \end{aligned}$$

by (76), which is equivalent to the third order determinant of equation (77).*

Exercise 12.—To show the product of two determinants as a determinant of the same order.

Let $p_0 = \sum_0^2 l c$, $p_1 = \sum m c$, $p_2 = \sum n c$, $q_0 = \sum \lambda c$, $q_1 = \sum \mu c$, $q_2 = \sum \nu c$; then $p_0 p_1 p_2 = [l_0, m_1, n_2]$, $q_0 q_1 q_2 = [\lambda_0, \mu_1, \nu_2]$; also

$p_0 | q_0 = l_0 \lambda_0 + l_1 \lambda_1 + l_2 \lambda_2$, $p_1 | q_0 = m_0 \lambda_0 + m_1 \lambda_1 + m_2 \lambda_2$, etc. Substituting these values in (77), we have the required result. A solution may also be obtained directly without the use of (77).

Let the q 's be as above, but write $p_0 = \sum_0^2 l q$, $p_1 = \sum m q$, $p_2 = \sum n q$.

Then

$$p_0 p_1 p_2 = \sum l q \cdot \sum m q \cdot \sum n q = [l_0, m_1, n_2] q_0 q_1 q_2 = [l_0, m_1, n_2] [\lambda_0, \mu_1, \nu_2].$$

* Grassmann (1862), Art. 173.

Also $p_0 = l_0 \sum \lambda e + l_1 \sum \mu e + l_2 \sum \nu e$

$$= (l_0 \lambda_0 + l_1 \mu_0 + l_2 \nu_0) e_0 + (l_0 \lambda_1 + l_1 \mu_1 + l_2 \nu_1) e_1 + (l_0 \lambda_2 + l_1 \mu_2 + l_2 \nu_2) e_2,$$

with similar values for p_1 and p_2 , which on being substituted in $p_0 p_1 p_2$ give the result. Equation (77), however, exhibits the product in a very compact, symmetrical, and easily remembered form.*


Exercise 13.—Show that the sides $p_1 p_2$, $p_2 p_3$, $p_3 p_1$ of the triangle $p_1 p_2 p_3$ cut the corresponding sides $|p_3|$, $|p_1|$, $|p_2|$ of the complementary triangle in three collinear points.

The three points of intersection are, using (74),

$p_1 p_2 \cdot |p_3| = -p_1 \cdot p_2 |p_3| + p_2 \cdot p_1 |p_3|$, $p_2 p_3 \cdot |p_1| = -p_2 \cdot p_3 |p_1| + p_3 \cdot p_2 |p_1|$, $p_3 p_1 \cdot |p_2| = -p_3 \cdot p_1 |p_2| + p_1 \cdot p_3 |p_2|$, of which the sum is zero, showing that the points are collinear. It may be shown in the same way that the lines joining corresponding vertices are confluent.

Exercise 14.—If the sides of a triangle pass through three fixed points, and two of the vertices slide on fixed lines, find the locus of the other vertex.

Let the fixed points and lines be p_1 , p_2 , p_3 , L_1 , L_2 , and p , p' , p'' the vertices of the triangle, as in the figure. Then $p' p_3 p'' = 0$; p' coincides with $p p_1 \cdot L_1$ and p'' with $p p_2 \cdot L_2$; hence substituting $(p p_1 \cdot L_1) p_3 (L_2 \cdot p_2 p) = 0$, the equation of the locus, which, being of the second degree in p , is that of a conic.



Prob. 18. Show that if the three fixed points of the last exercise are collinear, then the locus of p breaks up into two straight lines. Use equation (73).

Prob. 19. If the vertices of a triangle slide on three fixed lines, and two of the sides pass through fixed points, find the envelope of the other side. (This statement is reciprocally related to that of Exercise 14, that is, lines and points are replaced by points and

* These methods may be applied to determinants of any order by using a space of corresponding order.

lines respectively, and the resulting equation will be an equation of the second order in L , a variable line.)


Prob. 20. Show that if the three fixed lines of Problem 19 are confluent, then the envelope of L reduces to two points and the line joining them.

ART. 10. STEREOMETRIC PRODUCTS.

The product of two points in solid space is the same as in plane space. See Art. 7.

Product of Three Points.—Any three points determine a plane, and also, as in Art. 7, an area; hence $p_1 p_2 p_3$ is a plane-sect or a portion of the plane fixed by the three points whose area is double that of the triangle $p_1 p_2 p_3$. It may be shown, in the manner used in Art. 7 for the sect, that no plane-sect, not in this plane, can be equal to $p_1 p_2 p_3$, and that any plane-sect in this plane having the same area and sign will be equal to $p_1 p_2 p_3$.^{*} Of course $p_1 p_2 p_3$ is not now scalar.

Product of Four Points.—Any four non-coplanar points



determine a tetrahedron, say $p_1 p_2 p_3 p_4$, and six times the volume of this tetrahedron is taken for the value of the product, because this is the volume of the parallelepiped

generated by the product $p_1 p_2 p_3$,—i.e. the parallelogram p_1, p_3 ,—when it moves parallel to its initial position from p_1 to p_4 . Let $p_2 - p_1 = \epsilon$, $p_3 - p_1 = \epsilon'$, $p_4 - p_1 = \epsilon''$, then

$$p_1 p_2 p_3 p_4 = p_1 p_2 p_3 \epsilon'' = p_1 p_2 \epsilon' \epsilon'' = p_1 \epsilon \epsilon' \epsilon''. \quad (78)$$

If $p_1 = \sum_0^3 k e$, $p_2 = \sum_0^3 l e$, $p_3 = \sum_0^3 m e$, $p_4 = \sum_0^3 n e$, then

$$p_1 p_2 p_3 p_4 = \sum k e \sum l e \sum m e \sum n e = [k_0, l_1, m_2, n_3] \cdot e_0 e_1 e_2 e_3; \quad (79)$$

from which it appears that any two quadruple products of points differ from each other only by a scalar factor, that is, they differ only in magnitude, or sign, or both; hence such products are themselves scalar.[†] If $p_1 p_2 p_3 p_4 = 0$, the volume of the tetrahedron vanishes, so that the four points are coplanar.

^{*} Grassmann (1862), Art. 255.

[†] Grassmann (1862), Art. 263.

Product of Two Vectors.—The two vectors determine an area as in Art. 7, but they also determine now a plane direction, so that the product $\epsilon_1\epsilon_2$ is a plane-vector, and is not scalar as in plane space. Also, $\epsilon_1\epsilon_2$ differs from $p_1\epsilon_1\epsilon_2$ now just as ϵ differs from $p\epsilon$; namely, $\epsilon_1\epsilon_2$ has a definite area and plane direction, that is, toward a certain line at infinity, while $p_1\epsilon_1\epsilon_2$ is fixed in position by passing through p_1 . Equation (37) therefore does not hold in solid space.

Product of Three Vectors.—Three vectors determine a parallelepiped as in the figure above, and $\epsilon\epsilon'\epsilon''$ is therefore the volume of this parallelepiped. Any other triple vector product can differ from this only in magnitude and sign. For let $\epsilon_1\epsilon_2\epsilon_3$ be such a product, and write

$$\epsilon = x_1\epsilon_1 + x_2\epsilon_2 + x_3\epsilon_3 = \sum_1^3 x\epsilon, \quad \epsilon' = \sum_1^3 y\epsilon, \quad \epsilon'' = \sum_1^3 z\epsilon; \text{ then}$$

$$\epsilon\epsilon'\epsilon'' = \sum x\epsilon \sum y\epsilon \sum z\epsilon = \begin{vmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{vmatrix} \epsilon_1\epsilon_2\epsilon_3, \quad (80)$$

so that the two products only differ by the scalar determinant factor. Hence the product of three vectors must be itself a scalar, by Art. 1. Since, then, the product of four points has precisely the same signification as that of three vectors, we may write

$$\begin{aligned} p_1p_2p_3p_4 &= p_1\epsilon\epsilon'\epsilon'' = \epsilon\epsilon'\epsilon'' = (p_2 - p_1)(p_3 - p_1)(p_4 - p_1) \\ &= p_2p_3p_4 - p_3p_4p_1 + p_4p_1p_2 - p_1p_2p_3. \end{aligned} \quad (81)$$

Thus the sum of the plane-sects forming the doubles of the faces of a tetrahedron, all taken positively in the same sense as looked at from outside the tetrahedron, is equal to the volume of the tetrahedron. Compare equation (37).

If $\epsilon\epsilon'\epsilon'' = 0$, the volume of the parallelepiped vanishes, and the three vectors must be parallel to one plane.

Product of Two Sects.—In solid space two sects determine a tetrahedron of which they are opposite edges. Thus

$$p_1p_2p_3p_4 = p_1p_2 \cdot p_3p_4 = L_1L_2 = p_3p_4 \cdot p_1p_2 = L_2L_1, \quad (82)$$

so that the stereometric product of two sects is commutative, and has the same meaning as that of four points.

Product of a Sect and a Plane-Sect.—Let them be L and P , and let p_0 be their common point; take p_1, p_2, p_3 so that $L = p_0 p_1$ and $P = p_0 p_2 p_3$. L and P evidently determine the point p_0 , and also the parallelepiped of which one edge is L and one face is P , so that the product should be made up of these two factors. Hence we write

$$\left. \begin{aligned} LP &= p_0 p_1 \cdot p_0 p_2 p_3 = p_0 p_1 p_2 p_3 \cdot p_0; \\ PL &= p_0 p_2 p_3 \cdot p_0 p_1 = p_0 p_2 p_3 p_1 \cdot p_0 = LP. \end{aligned} \right\} \quad (83)$$

If L is parallel to P , p_0 is at infinity, and, replacing it by ϵ , (83) becomes

$$PL = LP = \epsilon p_1 \cdot \epsilon p_2 p_3 = \epsilon p_1 p_2 p_3 \cdot \epsilon. \quad (84)$$

Product of Two Plane-Sects.—Let them be P_1 and P_2 , and let L be their intersection, while p_1 and p_2 are such points that $P_1 = L p_1$ and $P_2 = L p_2$; then P_1 and P_2 determine the line L and also a parallelepiped of which they are two adjacent faces, and

$$P_1 P_2 = L p_1 \cdot L p_2 = L p_1 p_2 \cdot L = -P_2 P_1. \quad (85)$$

If P_1 and P_2 are parallel, L is at infinity, and is equivalent to a plane-vector, say to η ; hence, substituting in (84),

$$P_1 P_2 = \eta p_1 \cdot \eta p_2 = \eta p_1 p_2 \cdot \eta = -P_2 P_1. \quad (86)$$

Product of Three Plane-Sects.—By (85) and (83) this must be the square of a volume times the common point of the three planes; or, if p_0, p_1, p_2, p_3 be taken in such manner that $P_1 = p_0 p_2 p_3$, $P_2 = p_0 p_3 p_1$, $P_3 = p_0 p_1 p_2$, then

$$P_1 P_2 P_3 = 023 \cdot 031 \cdot 012 = 023 \cdot 0123 \cdot 01 = (p_0 p_1 p_2 p_3)^2 \cdot p_0; \quad (87)$$

the suffixes being used instead of the corresponding points. If p_0 be at infinity, the three planes are parallel to a single line, and may be written $P_1 = n_1 \epsilon p_2 p_3$, etc., and then treated as above.

Product of Four Plane-Sects.*—Let the planes be $P_0 \dots P_3$, and let $p_0 \dots p_3$ be the four common points of the planes taken three by three. $n_0 \dots n_3$ may be so taken that $P_0 = n_0 p_1 p_2 p_3$, etc.; then

$$\begin{aligned} P_0 P_1 P_2 P_3 &= n_0 n_1 n_2 n_3 \cdot 123 \cdot 230 \cdot 301 \cdot 012 \\ &= n_0 n_1 n_2 n_3 (p_0 p_1 p_2 p_3)^2. \end{aligned} \quad (88)$$

* Grassmann (1862), Art. 300.

Product of Two Plane-Vectors.—Let η_1 and η_2 be two plane-vectors or lines at infinity; let ϵ be parallel to each of them, and ϵ_1 and ϵ_2 so taken that $\eta_1 = \epsilon\epsilon_1$, $\eta_2 = \epsilon\epsilon_2$, then

$$\eta_1\eta_2 = \epsilon\epsilon_1 \cdot \epsilon\epsilon_2 = \epsilon\epsilon_1\epsilon_2 \cdot \epsilon = -\eta_2\eta_1, \quad (89)$$

because η_1 and η_2 determine a common direction ϵ , and a parallelepiped of which three conterminous edges are equal to ϵ , ϵ_1 , ϵ_2 , respectively.

Product of Three Plane-Vectors.—Take ϵ_1 , ϵ_2 , ϵ_3 so that

$$\eta_1\eta_2\eta_3 = n \cdot \epsilon_2\epsilon_3 \cdot \epsilon_3\epsilon_1 \cdot \epsilon_1\epsilon_2 = n(\epsilon_1\epsilon_2\epsilon_3)^2. \quad (90)$$

The directions $\epsilon_1 \dots \epsilon_3$ are common to the plane-vectors $\eta_1 \dots \eta_3$ taken two by two.

Several conditions are given here together which follow from the results of this article.

$$\rho_1\rho_2 = 0, \quad P_1P_2 = 0, \quad (91)$$

Two points coincide.

Two planes coincide.

$$\rho_1\rho_2\rho_3 = 0, \quad P_1P_2P_3 = 0, \quad (92)$$

Three points collinear.

Three planes collinear.

$$\begin{aligned} \rho_1\rho_2\rho_3\rho_4 &= \rho_1\rho_2 \cdot \rho_3\rho_4 \\ &= L_1L_2 = 0, \end{aligned} \quad \begin{aligned} P_1P_2P_3P_4 &= P_1P_2 \cdot P_3P_4 \\ &= L_1L_2 = 0, \end{aligned} \quad (93)$$

Four points coplanar; two lines intersect.

Four planes confluent; two lines intersect.

$$\epsilon_1\epsilon_2 = 0, \quad \eta_1\eta_2 = 0, \quad (94)$$

Vectors parallel.

Plane-vectors parallel.

$$\epsilon_1\epsilon_2\epsilon_3 = 0, \quad \eta_1\eta_2\eta_3 = 0, \quad (95)$$

Three vectors parallel to one plane.

Three plane-vectors parallel to one line.

Sum of Two Planes.—Let them be P_1 and P_2 , let L be a sect in their common line, and take ρ_1 and ρ_2 so that $P_1 = L\rho_1$, $P_2 = L\rho_2$; then

$$P_1 + P_2 = L(\rho_1 + \rho_2) = 2L\bar{\rho}, \quad (96)$$

$\bar{\rho}$ being the mean of ρ_1 and ρ_2 . Also

$$P_1 - P_2 = L(\rho_1 - \rho_2); \quad (97)$$

whence the sum and difference are the diagonal plane through L , and a plane through L parallel to the diagonal plane which is itself parallel to L , of the parallelepiped determined by P_1

and P_2 . If $TP_1 = TP_2$, $P_1 \pm P_2$ will evidently be the two bisecting planes of the angle between them. The bisecting planes may also be written

$$\frac{P_1}{TP_1} \pm \frac{P_2}{TP_2} \quad \text{or} \quad P_1 TP_2 \pm P_2 TP_1. \quad (98)$$

If the two planes are parallel, let η be a plane-vector parallel to each of them, that is, their common line at infinity, and let p_1 and p_2 be points in the respective planes; then we may write $P_1 = n_1 p_1 \eta$, $P_2 = n_2 p_2 \eta$, whence

$$P_1 + P_2 = (n_1 p_1 + n_2 p_2) \eta = (n_1 + n_2) \bar{p} \eta. \quad (99)$$

If $n_1 + n_2 = 0$, this becomes

$$P_1 + P_2 = n_2 (p_2 - p_1) \eta, \quad (100)$$

the product of a vector into a plane-vector and therefore a scalar, by (80).

Two plane-vectors may be added similarly, since they will have a common direction, namely, that of the vector parallel to both of them.

Exercise 15.—If two tetrahedra $e_0 e_1 e_2 e_3$ and $e'_0 e'_1 e'_2 e'_3$ are so situated that the right lines through the pairs of corresponding vertices all meet in one point, then will the corresponding faces cut each other in four coplanar lines.

The given conditions are equivalent to $e_0 e'_0 \cdot e_1 e'_1 = 0 = e_0 e'_0 \cdot e_2 e'_2 = e_0 e'_0 \cdot e_3 e'_3 = e_1 e'_1 \cdot e_2 e'_2 = e_1 e'_1 \cdot e_3 e'_3 = e_2 e'_2 \cdot e_3 e'_3 = e_2 e'_2 \cdot e_1 e'_1$. Two of the intersecting lines of faces are $e_0 e_1 e_2 \cdot e'_0 e'_1 e'_2$ and $e_1 e_2 e_3 \cdot e'_1 e'_2 e'_3$, and, if these intersect, we must accordingly have, by (93), $012 \cdot 0'1'2' \cdot 123 \cdot 1'2'3' = 0 = 012 \cdot 123 \cdot 0'1'2' \cdot 1'2'3' = 0123 \cdot 0'1'2'3' \cdot 121'2'$, the last factor of which is equivalent to the fourth condition above, since quadruple-point products in solid space are associative. Similarly all the other pairs of intersections may be treated.

Exercise 16.—The twelve bisecting planes of the diedral angles of a tetrahedron fix eight points, the centers of the inscribed and escribed spheres, through which they pass six by six.

The sum and difference of two unit planes are their two

bisecting planes, by (97). Let the tetrahedron be $e_0e_1e_2e_3$, and let the double areas of its faces be $A_0 = Te_1e_2e_3$, etc.; then a pair of bisecting planes will be $\frac{e_0e_1e_2}{A_3} \pm \frac{e_0e_1e_3}{A_2}$ or $e_0e_1(A_2e_2 \pm A_3e_3)$. The pair through the opposite edge will be $e_2e_3(A_0e_0 \pm A_1e_1)$. If there be a point through which the six internal bisecting planes pass, it must be on the intersection of these two planes taken with the upper signs, and we infer by symmetry that it must be the point $\sum_0^3 Ae$. Another internal bisecting plane is $e_3e_0(A_1e_1 + A_2e_2)$, which gives zero when multiplied into $\sum Ae$, as do also the other three.

To obtain all the points we have only to use the double signs, so that they are $\pm A_0e_0 \pm A_1e_1 \pm A_2e_2 \pm A_3e_3$. This gives eight cases, namely,

+ + + +	- + + +
+ + + -	+ + - -
+ + - +	+ - - +
+ - + +	+ - + -

The eight apparent cases that would arise by changing all the signs are included in these because the points must be essentially positive. Moreover, no positive point could have three negative signs, because the sum of any three faces of the tetrahedron must be greater than the fourth face. It will be found on trial that six of the bisecting planes will pass through $\sum(\pm Ae)$ with any one of the above arrangements of sign.

Prob. 21. The twelve points in which the edges of a tetrahedron are cut by the bisecting planes of the opposite dihedral angles fix eight planes, each of which passes through six of them.

Prob. 22. The centroid of the faces of a tetrahedron coincides with the center of the sphere inscribed within the tetrahedron whose vertices are the centroids of the respective faces of the first tetrahedron.

Prob. 23. If any plane be passed through the middle points of two opposite edges of a tetrahedron, it will divide the volume of the tetrahedron into two equal parts.

ART. 11. THE COMPLEMENT IN SOLID SPACE.


According to the definitions of Art. 8 the complementary relations in a unit normal vector system are as follows:

$$\left. \begin{aligned} |l_1 = l_2 l_3, & \quad |l_2 l_3 = |(|l_1) = l_1 \\ |l_2 = l_3 l_1, & \quad |l_3 l_1 = |(|l_2) = l_2 \\ |l_3 = l_1 l_2, & \quad |l_1 l_2 = |(|l_3) = l_3 \end{aligned} \right\}. \quad (101)$$

Let $\epsilon = \sum_1^3 l_i$; then

$$|\epsilon = l_1 l_2 l_3 + l_2 l_3 l_1 + l_3 l_1 l_2 = \frac{1}{l_1} (l_1 l_2 - l_2 l_1) (l_1 l_3 - l_3 l_1), \quad (102)$$

so that $|\epsilon$ is a plane-vector. The figure, which is drawn in



isometric projection, shows that the two vectors $l_1 l_2 - l_2 l_1$ and $l_1 l_3 - l_3 l_1$, whose product is $l_1 \cdot |\epsilon$, are both perpendicular to ϵ ; for the first is perpendicular to $l_1 l_1 + l_2 l_2$, which is the orthogonal projection of ϵ upon $l_1 l_2$, and to l_3 , and therefore is also perpendicular to $\bar{\epsilon}$, while the

second is perpendicular to $l_1 l_1 + l_3 l_3$ and to l_2 , and therefore to ϵ . Hence $|\epsilon$ is a plane-vector perpendicular to ϵ ; and, since $|(|\epsilon) = \epsilon$, the converse is also true, i.e. the complement of a plane-vector is a line-vector normal to it.

The figure shows that ϵ is equal to the vector diagonal of the rectangular parallelepiped whose edges have the lengths l_1, l_2, l_3 , hence

$$T\epsilon = \sqrt{l_1^2 + l_2^2 + l_3^2}. \quad (103)$$

Multiply equation (102) by ϵ ; therefore

$$\begin{aligned} \epsilon |\epsilon &= (l_1 l_1 + l_2 l_2 + l_3 l_3) (l_1 l_2 l_3 + l_2 l_3 l_1 + l_3 l_1 l_2) \\ &= l_1^2 + l_2^2 + l_3^2 = T^2 \epsilon = \epsilon^2, \end{aligned} \quad (104)$$

so that the co-square of a vector is equal to the square of its tensor. The product $\epsilon |\epsilon$ is that of a vector ϵ into a plane-vector perpendicular to it, as has just been shown; it is there-


fore a volume which is equivalent to $T\epsilon \cdot T|\epsilon$; hence, by (104), $\epsilon|\epsilon = T\epsilon \cdot T|\epsilon = T^2\epsilon$, or $T\epsilon = T|\epsilon$. Hence, the complement of a vector in solid space is a plane-vector perpendicular to it and having the same tensor, or numerical measure of magnitude.*

Let a second vector be $\epsilon' = \sum_1^3 m_i$; then

$$\epsilon|\epsilon' = l_1 m_1 + l_2 m_2 + l_3 m_3 = \epsilon'|\epsilon. \quad (105)$$

Now $\epsilon|\epsilon'$ being the product of ϵ into the plane-vector $|\epsilon'$, is the volume of the parallelepiped in the figure, that is, $T\epsilon T\epsilon' \sin(\text{angle between } \epsilon \text{ and } |\epsilon') = T\epsilon T\epsilon' \cos \epsilon'$. Hence

$$\epsilon|\epsilon' = \epsilon'|\epsilon = l_1 m_1 + l_2 m_2 + l_3 m_3 = T\epsilon T\epsilon' \cos \epsilon'. \quad (106)$$



If $T\epsilon = T\epsilon' = 1$, $l_1 \dots l_3$, $m_1 \dots m_3$ are direction cosines, and (105) gives a proof of the formula for the cosine of the angle between two lines in terms of the direction cosines of the lines. We have also in this case

$\epsilon\epsilon' = (l_1 m_2 - l_2 m_1)|\iota_3 + (l_2 m_3 - l_3 m_2)|\iota_1 + (l_3 m_1 - l_1 m_3)|\iota_2$, and, taking the co-square,

$$(\epsilon\epsilon')^2 = (\sin \epsilon')^2 = (l_1 m_2 - l_2 m_1)^2 + (l_2 m_3 - l_3 m_2)^2 + (l_3 m_1 - l_1 m_3)^2. \quad (107)$$

$$\text{If} \quad \epsilon|\epsilon' = 0, \quad (108)$$

ϵ is parallel to the plane-vector perpendicular to ϵ' , that is, ϵ is perpendicular to ϵ' , as is also shown by (106).

Let $\eta = |\epsilon$, $\eta' = |\epsilon'$; then

$$\eta|\eta' = |\epsilon \cdot \epsilon' = \epsilon'|\epsilon = \epsilon|\epsilon' = T\epsilon T\epsilon' \cos \epsilon' = T\eta T\eta' \cos \eta', \quad (109)$$

$$\text{and} \quad \eta|\eta' = 0 \quad (110)$$

is the condition of perpendicularity of two plane-vectors. Also either

$$\epsilon|\eta' = 0, \quad \text{or} \quad \eta'|\epsilon = 0, \quad (111)$$

is the condition that a vector shall be perpendicular to a plane-vector, for the first means that ϵ is parallel to a vector which is

* Grassmann (1862), Art. 335.

perpendicular to η' , and the second that η' is parallel to a plane-vector which is perpendicular to ϵ .

Equations (71)–(77) of Art. 9 become stereometric vector formulæ if ϵ_1, ϵ_2 , etc., be substituted for p_1, p_2 , etc., and η_1, η_2 etc., for L_1, L_2 , etc. For instance, (76) gives the vector formulas

$$\epsilon_1 \epsilon_2 | \epsilon_1' \epsilon_2' = \begin{vmatrix} \epsilon_1 | \epsilon_1' & \epsilon_1 | \epsilon_2' \\ \epsilon_2 | \epsilon_1' & \epsilon_2 | \epsilon_2' \end{vmatrix}, \quad \eta_1 \eta_2 | \eta_1' \eta_2' = \begin{vmatrix} \eta_1 | \eta_1' & \eta_1 | \eta_2' \\ \eta_2 | \eta_1' & \eta_2 | \eta_2' \end{vmatrix}. \quad (112)$$

For lack of space no treatment of the complement in a point system in solid space is given.

Exercise 17.—To prove the formulas of spherical trigonometry $\cos a = \cos b \cos c + \sin b \sin c \cos A$, and

$$\frac{\sin a}{\sin A} = \frac{\sin b}{\sin B} = \frac{\sin c}{\sin C}.$$

Take three unit vectors $\epsilon_1, \epsilon_2, \epsilon_3$ parallel to the radii to the vertices of the spherical triangle, then $a = (\text{angle bet. } \epsilon_2 \text{ and } \epsilon_3)$, $A = (\text{angle bet. } \epsilon_1 \epsilon_2 \text{ and } \epsilon_1 \epsilon_3)$, etc. In eq. (112) put $\epsilon_1 \epsilon_3$ for $\epsilon_1' \epsilon_2'$;

$$\begin{aligned} \text{hence } \epsilon_1 \epsilon_2 | \epsilon_1 \epsilon_3 &= \sin b \sin c \cos A = \epsilon_2 | \epsilon_3 - \epsilon_1 | \epsilon_2 \cdot \epsilon_1 | \epsilon_3 \\ &= \cos a - \cos b \cos c. \end{aligned}$$

Again,

$$T(\epsilon_1 \epsilon_2 \cdot \epsilon_1 \epsilon_3) = T(\epsilon_1 \epsilon_2 \epsilon_3 \cdot \epsilon_1) = T\epsilon_1 \epsilon_2 \epsilon_3 = T(\epsilon_2 \epsilon_3 \cdot \epsilon_2 \epsilon_1) = T(\epsilon_3 \epsilon_1 \cdot \epsilon_3 \epsilon_2);$$

$$\text{or } \sin b \sin c \sin A = \sin a \sin c \sin B = \sin a \sin b \sin C,$$

whence we have the second result by dividing by $\sin a \sin b \sin c$.

Exercise 18.—Show that in a spherical triangle taken as in Exercise 17, $\cos \frac{A}{2} = \frac{U\epsilon_1 \epsilon_2 | (U\epsilon_1 \epsilon_2 + U\epsilon_1 \epsilon_3)}{T(U\epsilon_1 \epsilon_2 + U\epsilon_1 \epsilon_3)}$, whence derive

$$\text{the ordinary value } \sqrt{\frac{\sin s \sin (s - a)}{\sin b \sin c}}.$$

Expanding, the numerator becomes $1 + U\epsilon_1 \epsilon_2 | U\epsilon_1 \epsilon_3$, and the denominator $\sqrt{2(1 + U\epsilon_1 \epsilon_2 | U\epsilon_1 \epsilon_3)}$. Also there is obtained

$$U\epsilon_1 \epsilon_2 | U\epsilon_1 \epsilon_3 = \frac{\epsilon_1 \epsilon_2 \cdot \epsilon_1 \epsilon_3}{T\epsilon_1 \epsilon_2 T\epsilon_1 \epsilon_3}.$$

The remainder is left to the student.

Prob. 24. If $\epsilon_1, \epsilon_2, \epsilon_3$, drawn outward from a point, are taken as three edges of a tetrahedron, show that the six planes perpen-

dicular to the edges at their middle points all pass through the end of the vector $\rho = \frac{1}{2\epsilon_1\epsilon_2\epsilon_3}(|\epsilon_2\epsilon_3 \cdot \epsilon_1|^2 + |\epsilon_3\epsilon_1 \cdot \epsilon_2|^2 + |\epsilon_1\epsilon_2 \cdot \epsilon_3|^2)$. (Suggestion. We must have $(\rho - \frac{1}{2}\epsilon_1)|\epsilon_1| = 0$, with two other similar expressions.)

Prob. 25. Show that ϵ , $|\epsilon\epsilon'$ and $\epsilon\epsilon'$, $|\epsilon$ are three mutually perpendicular vectors, no matter what the directions of ϵ and ϵ' may be.

Prob. 26. Let $\epsilon_1, \epsilon_2, \epsilon_3$ be taken as in Prob. 24; let A_0 be the area of the face of the tetrahedron formed by joining the ends of these vectors, and $2A_1 = T\epsilon_2\epsilon_3$, etc.; also $\theta_1 = \text{Angle between } \epsilon_1\epsilon_2 \text{ and } \epsilon_1\epsilon_3$, etc.: then show that we have the relation, analogous to that of Prob. 15, Art. 8,

$A_0^2 = A_1^2 + A_2^2 + A_3^2 - 2A_2A_3 \cos \theta_1 - 2A_3A_1 \cos \theta_2 - 2A_1A_2 \cos \theta_3$. If $\theta_1 \dots \theta_3$ are right angles, this becomes the space-analog of the proposition regarding the hypotenuse and sides of a right-angled triangle. (Suggestion. $2A_0 = T(\epsilon_2 - \epsilon_1)(\epsilon_3 - \epsilon_1)$.)

Prob. 27. There are given three non-coplanar lines $\epsilon_0\epsilon_1, \epsilon_0\epsilon_2, \epsilon_0\epsilon_3$; planes cut these lines at right angles, the sum of the squares of their distances from ϵ_0 being constant. Show that the locus of the common point of these three planes is $(\rho|\epsilon_1|^2 + (\rho|\epsilon_2|^2 + (\rho|\epsilon_3|^2 = c^2$, if $T\epsilon_1 = T\epsilon_2 = T\epsilon_3 = 1$.

ART. 12. ADDITION OF SECTS IN SOLID SPACE.

Two lines in solid space will not in general intersect, so that their sum will not be, as in eq. (43), a definite line. For let $p_1\epsilon_1$ and $p_2\epsilon_2$ be any two sects: then

$$\begin{aligned} p_1\epsilon_1 + p_2\epsilon_2 &= p_1\epsilon_1 + p_2\epsilon_2 + \epsilon_0(\epsilon_1 + \epsilon_2) - \epsilon_0(\epsilon_1 + \epsilon_2) \\ &= \epsilon_0(\epsilon_1 + \epsilon_2) + (p_1 - \epsilon_0)\epsilon_1 + (p_2 - \epsilon_0)\epsilon_2; \end{aligned}$$

that is, the sum is a sect passing through an arbitrary point ϵ_0 , and a plane-vector, the sum of the two in the equation. The sum cannot be a single sect unless the two are coplanar; for let $p_2 = p_1 + x\epsilon_1 + y\epsilon_2 + z\epsilon_3$, ϵ_3 being a vector not parallel to $\epsilon_1\epsilon_2$;

$$\begin{aligned} p_1\epsilon_1 + p_2\epsilon_2 &= p_1\epsilon_1 + (p_1 + x\epsilon_1 + y\epsilon_2 + z\epsilon_3)\epsilon_2 \\ &= p_1(\epsilon_1 + \epsilon_2) + x\epsilon_1(\epsilon_1 + \epsilon_2) + z\epsilon_3\epsilon_2 \\ &= (p_1 + x\epsilon_1)(\epsilon_1 + \epsilon_2) + z\epsilon_3\epsilon_2; \end{aligned}$$

and this cannot reduce to a single sect unless $z = 0$, that is, unless $p_1\epsilon_1$ and $p_2\epsilon_2$ are coplanar. Since a plane-vector is a line at

∞ , the sum of two lines may always be presented as the sum of a finite line and a line at ∞ .

If the sum of any two sects is equal to the sum of any other two, their products will also be equal, that is, the two pairs will determine tetrahedra of equal volumes. For let $L_1 + L_2 = L_3 + L_4$; then squaring we have $L_1 L_2 = L_3 L_4$, since $L_1 L_1 = 0$, etc.

An infinite number of pairs of sects can be found such that the sum of each pair is equal to the sum of any given pair; for let a given pair be $p_1 \epsilon_1 + p_2 \epsilon_2$, and take a new pair

$$\begin{aligned} & (x_1 p_1 + x_2 p_2)(u_1 \epsilon_1 + u_2 \epsilon_2) + (y_1 p_1 + y_2 p_2)(v_1 \epsilon_1 + v_2 \epsilon_2) \\ &= (x_1 u_1 + y_1 v_1) p_1 \epsilon_1 + (x_2 u_2 + y_2 v_2) p_2 \epsilon_2 + \\ & \quad (x_1 u_2 + y_1 v_2) p_1 \epsilon_2 + (x_2 u_1 + y_2 v_1) p_2 \epsilon_1. \end{aligned}$$

This will be equal to the given pair if we have

$$x_1 u_1 + y_1 v_1 = x_2 u_2 + y_2 v_2 = 1, \text{ and } x_1 u_2 + y_1 v_2 = x_2 u_1 + y_2 v_1 = 0.$$

Since there are eight arbitrary quantities with only four equations of condition, the desired result can evidently be accomplished in an infinite number of ways.

Let $p_1 \epsilon_1, p_2 \epsilon_2, \dots, p_n \epsilon_n$ be n sects, and let S be their sum, and e_0 any point, then

$$S = \sum_1^n p \epsilon \equiv e_0 \sum \epsilon - e_0 \sum \epsilon + \sum p \epsilon = e_0 \sum \epsilon + \sum (p - e_0) \epsilon, \dots \quad (113)$$

the sum of a sect and a plane-vector as before.

If $\sum (p - e_0) \epsilon$ is parallel to $\sum \epsilon$ it may be written as the product of some vector ϵ' into $\sum \epsilon$, that is, $\epsilon' \sum \epsilon$, when the sum becomes $S = e_0 \sum \epsilon + \epsilon' \sum \epsilon = (e_0 + \epsilon') \sum \epsilon$, a sect, because $e_0 + \epsilon'$ is a point. In no other case does S reduce to a single sect. If $\sum \epsilon = 0$ S becomes a plane-vector. Of the two parts composing S , the sect will be unchanged in magnitude and direction if e_0 be moved to a new position, while the plane-vector will in general be altered. It is proposed to show that a point q may be substituted for e_0 such that the plane-vector will be perpendicular to $\sum \epsilon$. Writing

$$S \equiv q \sum \epsilon - (q - e_0) \sum \epsilon + \sum (p - e_0) \epsilon,$$

and, for brevity, putting $q - e_0 = \rho$, $\sum \epsilon = \alpha$, $\sum (p - e_0) \epsilon = |\beta$, so that

$$S \equiv q \alpha - \rho \alpha + |\beta, \quad (114)$$

we must have for perpendicularity, by (111),

$$\begin{aligned} &(|\beta - \rho\alpha| \mid \alpha = 0 = |\beta\alpha - \rho\alpha \cdot | \alpha, \\ \text{or} \quad &\rho\alpha \cdot | \alpha \equiv \alpha \cdot \rho \mid \alpha - \rho \cdot \alpha^2 = |\beta\alpha. \end{aligned} \quad (115)$$

The second member is obtained from the first by substituting in eq. (74) ρ for p_1 and α for p_2 and q_1 , in accordance with the statement at the end of Art. 11. If in (115) we make $\rho \mid \alpha = 0$, ρ will be the vector from e_0 to q taken perpendicularly to α , say

$$\rho_1 = |\alpha\beta \div \alpha^2 = q_1 - e_0. \quad (116)$$

Since α and β are known, the required point has been found. Multiply (115) by α ; then, using (75),

$$- \alpha\rho \cdot \alpha^2 \equiv \rho\alpha \cdot \alpha^2 = \alpha \mid \beta\alpha = |\beta \cdot \alpha^2 - |\alpha \cdot \alpha \mid \beta,$$

whence, substituting in (114),

$$S = q\alpha + \frac{\alpha \mid \beta}{\alpha^2} \cdot | \alpha = q\Sigma\epsilon + \frac{\Sigma\epsilon\Sigma(p - e_0)\epsilon}{(\Sigma\epsilon)^2} \cdot | \Sigma\epsilon. \quad (117)$$

This may be called the normal form of S .*

The sects of this article represent completely the geometric properties of forces, hence all that has been shown applies immediately to a system of forces in solid space. We have only to substitute the words force and couple for sect and plane-vector. The resultant action of any system of forces is S , called by Ball in his Theory of Screws "a wrench." The condition for equilibrium is $S = 0$, which gives at once

$$\Sigma\epsilon = 0 \quad \text{and} \quad \Sigma(p - e_0)\epsilon = 0; \quad (118)$$

since otherwise we must have $e_0\Sigma\epsilon = -\Sigma(p - e_0)\epsilon$, which is an impossibility. The line $q\Sigma\epsilon$ is the central axis of the system of forces S .

Lack of space forbids a further development of the subject, but what has been given in this article will indicate the perfect adaptability of this method to the requirements of mechanics.

Exercise 19.—Reduce $p_1\epsilon_1 + p_2\epsilon_2 = S$ to its normal form. $S \equiv e_0(\epsilon_1 + \epsilon_2) + (p_1 - e_0)\epsilon_1 + (p_2 - e_0)\epsilon_2$. For convenience suppose p_1 and p_2 to be taken at the ends of the common per-

* Grassmann (1862), Art. 346.

pendicular on $p_1\epsilon_1$ and $p_2\epsilon_2$, and moreover let $e_0 = \frac{1}{2}(p_1 + p_2)$, $p_1 - e_0 = \iota = -(p_2 - e_0)$; then $\iota|\epsilon_1 = \iota|\epsilon_2 = 0$. Accordingly

$$S \equiv e_0(\epsilon_1 + \epsilon_2) + \iota(\epsilon_1 - \epsilon_2) = q(\epsilon_1 + \epsilon_2) + \frac{(\epsilon_1 + \epsilon_2)\iota(\epsilon_1 - \epsilon_2)}{(\epsilon_1 + \epsilon_2)^2} \cdot |(\epsilon_1 + \epsilon_2)$$

$$= q(\epsilon_1 + \epsilon_2) + \frac{2\iota\epsilon_1\epsilon_2}{(\epsilon_1 + \epsilon_2)^2} \cdot |(\epsilon_1 + \epsilon_2).$$

$$\text{By (116), } q - e_0 = -\frac{|\beta \cdot | \alpha}{\alpha^2} = -\frac{\iota(\epsilon_1 - \epsilon_2) \cdot |(\epsilon_1 + \epsilon_2)}{(\epsilon_1 + \epsilon_2)^2}$$

$$= \frac{\iota \cdot (\epsilon_1 - \epsilon_2) |(\epsilon_1 + \epsilon_2)}{(\epsilon_1 + \epsilon_2)^2}, \text{ by (74), } = \frac{\epsilon_1^2 - \epsilon_2^2}{(\epsilon_1 + \epsilon_2)^2} \cdot \iota.$$

Hence the normal form of S is

$$S = \left(e_0 + \frac{\epsilon_1^2 - \epsilon_2^2}{(\epsilon_1 + \epsilon_2)^2} \cdot \iota \right) (\epsilon_1 + \epsilon_2) + \frac{2\iota\epsilon_1\epsilon_2}{(\epsilon_1 + \epsilon_2)^2} \cdot |(\epsilon_1 + \epsilon_2).$$

Exercise 20.—Forces are represented by the six edges of a tetrahedron $e_0e_1, e_0e_2, e_0e_3, e_2e_3, e_3e_1, e_1e_2$; find the S , reduce to normal form, and consider the special case when three diedral angles are right angles. $S \equiv e_0(e_1 + e_2 + e_3) + e_2e_3 + e_3e_1 + e_1e_2 \equiv e_0(\epsilon_1 + \epsilon_2 + \epsilon_3) + (\epsilon_2 - e_1)(\epsilon_3 - e_1) \equiv e_0(\epsilon_1 + \epsilon_2 + \epsilon_3) + (\epsilon_2 - e_1)(\epsilon_3 - e_1) \equiv e_0(\epsilon_1 + \epsilon_2 + \epsilon_3) + \epsilon_2\epsilon_3 + \epsilon_3\epsilon_1 + \epsilon_1\epsilon_2$, in which $\epsilon_1 = e_1 - e_0$, etc. Hence

$$S \equiv \left(e_0 + \frac{(\epsilon_2\epsilon_3 + \epsilon_3\epsilon_1 + \epsilon_1\epsilon_2) |(\epsilon_1 + \epsilon_2 + \epsilon_3)}{(\epsilon_1 + \epsilon_2 + \epsilon_3)^2} \right) (\epsilon_1 + \epsilon_2 + \epsilon_3)$$

$$+ \frac{3\epsilon_1\epsilon_2\epsilon_3}{(\epsilon_1 + \epsilon_2 + \epsilon_3)^2} \cdot |(\epsilon_1 + \epsilon_2 + \epsilon_3).$$

For the rectangular tetrahedron let $\epsilon_1 = a\iota_1, \epsilon_2 = b\iota_2, \epsilon_3 = c\iota_3, \iota_1, \iota_2, \iota_3$ being unit normal vectors. Then we find

$$S \equiv \left(e_0 + \frac{a(c^2 - b^2)\iota_1 + b(a^2 - c^2)\iota_2 + c(b^2 - a^2)\iota_3}{a^2 + b^2 + c^2} \right) (a\iota_1 + b\iota_2 + c\iota_3)$$

$$+ \frac{3abc}{a^2 + b^2 + c^2} \cdot | (a\iota_1 + b\iota_2 + c\iota_3).$$

Exercise 21.—A pole 50 feet high stands on the ground and is held erect by three guy-ropes symmetrically arranged about it, attached to its top and to pegs in the ground 50 feet from the pole. The wind blows against the pole with a pressure of 50 pounds in the direction $e_0 - p$, when e_0 is at the bottom of

the pole, and p divides the distance between two of the pegs in the ratio $\frac{m}{n}$: find the tension on the guys and the pressure on the ground.

Evidently only two of the guys will be in tension; let their pegs be at e_1 and e_3 , and let e_2 be at the top of the pole, and w the weight of the pole. Then $p = \frac{me_1 + ne_3}{m + n}$, and the equation of equilibrium is

$$50 \cdot \frac{(e_0 + e_2)(e_0 - p)}{2T(e_0 - p)} + \frac{25e_0(p - e_0)}{T(e_0 - p)} + \frac{(x + w)e_0e_2}{Te_0e_2} + \frac{ye_2e_1}{Te_2e_1} + \frac{ze_2e_3}{Te_2e_3} = 0.$$

$$Te_0e_2 = 50, Te_2e_1 = Te_2e_3 = 50\sqrt{2}, T(p - e_0) = T\left(\frac{me_1 + ne_3}{m + n} - e_0\right) \\ = T\left(\frac{m(e_1 - e_0) + n(e_3 - e_0)}{m + n}\right) = \frac{50}{m + n}T(m\epsilon_1 + n\epsilon_3), \text{ if } \epsilon_1 = U(e_1 - e_0)$$

$$\text{and } \epsilon_2 = U(e_2 - e_0); \text{ then } T(p - e_0) = \frac{50}{m + n}\sqrt{m^2 + n^2 - mn},$$

because $\epsilon_1^2 = \epsilon_3^2 = 1$, and $\epsilon_1\epsilon_3 = \cos 120^\circ = -\frac{1}{2}$. Hence the equation of equilibrium becomes

$$\frac{25e_2((m + n)e_0 - me_1 - ne_3)}{\sqrt{m^2 + n^2 - mn}} + (x + w)e_0e_2 + \frac{y}{\sqrt{2}}e_2e_1 + \frac{z}{\sqrt{2}}e_2e_3 = 0.$$

Multiply successively by e_3e_1 , e_0e_3 , and e_0e_1 , and we obtain

$$\frac{x + w}{m + n} = \frac{y}{m\sqrt{2}} = \frac{z}{n\sqrt{2}} = \frac{25}{\sqrt{m^2 + n^2 - mn}},$$

y and z being the tensions, and $x + w$ the upward pressure.

Prob. 28. Three equal poles are set up so as to form a tripod, and are mutually perpendicular; a weight w hangs upon a rope which passes over a pulley at the top of the tripod, and thence down under a pulley at the ground at a point $p = \sum_1^3 le$, in which $e_1 \dots e_3$ are at the feet of the poles, and $\sum_1^3 l = 1$; if the rope is pulled

so as to raise w , show that the pressures on the poles, supposing the pulleys frictionless, are

$$w\left(\frac{l_1}{\sqrt{\sum l^2}} + \frac{1}{\sqrt{3}}\right), \quad w\left(\frac{l_2}{\sqrt{\sum l^2}} + \frac{1}{\sqrt{3}}\right), \quad w\left(\frac{l_3}{\sqrt{\sum l^2}} + \frac{1}{\sqrt{3}}\right).$$

Prob. 29. Six equal forces act along six successive edges of a cube which do not meet a given diagonal; show that if the edges of the cube be parallel to $\iota_1, \iota_2, \iota_3$, and F be the magnitude of each force, then $S = -2F(\iota_1 + \iota_2 + \iota_3)$, if the diagonal taken be parallel to $\iota_1 + \iota_2 + \iota_3$.

Prob. 30. Three forces whose magnitudes are 1, 2, and 3 act along three successive non-coplanar edges of a cube; show that the normal form of S is

$$S = (e_0 + \frac{3}{14}\iota_1 + \frac{1}{2}\iota_2 - \frac{9}{14}\iota_3)(\iota_1 + 2\iota_2 + 3\iota_3) + \frac{3}{14}(\iota_1 + 2\iota_2 + 3\iota_3).$$

Prob. 31. Forces act at the centroids of the faces of a tetrahedron, perpendicular and proportional to the faces on which they act, and all directed inwards, or else all outwards; show that they are in equilibrium.

INDEX.

- Addition of points, page 9.
 of weighted points, 12-16.
 of vectors, 12.
 of sects (planimetric), 30.
 of sects (stereometric), 53.
 of plane-sects, 47.
- Coincidence of two points, 17, 39, 47.
- Collinearity of three points, 17, 39, 47.
- Coplanarity of four points, 17, 39, 47.
- Collinearity of ends of three vectors, 18.
- Coplanarity of ends of four vectors, 18.
- Combinatory products, 24.
 multiplication, Laws of, 26.
- Complement, 33, 50.
- Condition, equations of, 17, 39, 47.
- Co-product, 35.
- Co-square, 36.
- Difference of points, 10.
- Difference between p p_2 and $p_2 - p_1$, 24.
- Determinants, product, 42.
- Equations of condition, 17, 39, 47.
- Formulæ for points and lines in plain space, 41.
 for vectors in solid space, 52.
- Geometric multiplication, 24.
- Inner product, 35.
- Inscribed and escribed spheres, centers, 48.
- Laws of combinatory multiplication, 26.
- Mean point, 13.
- Multiplication, Geometric, 24.
 Combinatory, 24.
- Parallelism of vectors, 47.
 of plane-vectors, 47.
- Perpendicularity, Condition of, 36, 51.
- Plane sects, Product, 46.
 vectors, Product, 47.
 parallel, 47.
 three parallel to one line, 47.
- Planimetric products, 25, 26.
- Point at infinity = vector, 9.
- Product, Combinatory, 24.
- Products of two points, 26.
 of three points, 27.
 of two vectors, 28.
 of two sects, 28.
 of three sects, 29.
 of two determinants, 42.
- Reference systems, 20.
- Scalar, definition, 9.
- Sect, definition, 8.
- Sects, products, 28, 29, 45.
 products of parallel, 30.
 planimetric sum, 31.
 stereometric sum, 53.
- Spheres, inscribed and escribed, centers of, 48.
- Spherical trigonometry, formulæ, 52.
- Tensor = T , 9.
- Unit = U , 9.
- Vector, Definition of, 10.
 plane, 45.
- Vector products and conditions, 17, 18, 47.
- Wrench, 55.

SHORT-TITLE CATALOGUE

OF THE
PUBLICATIONS
OF
JOHN WILEY & SONS,
NEW YORK.

LONDON: CHAPMAN & HALL, LIMITED.

ARRANGED UNDER SUBJECTS.

Descriptive circulars sent on application. Books marked with an asterisk (*) are sold at *net* prices only, a double asterisk (**) books sold under the rules of the American Publishers' Association at *net* prices subject to an extra charge for postage. All books are bound in cloth unless otherwise stated.

AGRICULTURE.

Armsby's Manual of Cattle-feeding.	12mo,	\$1 75
Principles of Animal Nutrition.	8vo,	4 00
Budd and Hansen's American Horticultural Manual:		
Part I. Propagation, Culture, and Improvement.	12mo,	1 50
Part II. Systematic Pomology.	12mo,	1 50
Downing's Fruits and Fruit-trees of America.	8vo,	5 00
Elliott's Engineering for Land Drainage.	12mo,	1 50
Practical Farm Drainage.	12mo,	1 00
Green's Principles of American Forestry.	12mo,	1 50
Grotenfelt's Principles of Modern Dairy Practice. (Woll.).	12mo,	2 00
Kemp's Landscape Gardening.	12mo,	2 50
Maynard's Landscape Gardening as Applied to Home Decoration.	12mo,	1 50
* McKay and Larsen's Principles and Practice of Butter-making.	8vo,	1 50
Sanderson's Insects Injurious to Staple Crops.	12mo,	1 50
Insects Injurious to Garden Crops. (In preparation.)		
Insects Injuring Fruits. (In preparation.)		
Stockbridge's Rocks and Soils.	8vo,	2 50
Winton's Microscopy of Vegetable Foods.	8vo,	7 50
Woll's Handbook for Farmers and Dairymen.	16mo,	1 50

ARCHITECTURE.

Baldwin's Steam Heating for Buildings.	12mo,	2 50
Bashore's Sanitation of a Country House.	12mo,	1 00
Berg's Buildings and Structures of American Railroads.	4to,	5 00
Birkmire's Planning and Construction of American Theatres.	8vo,	3 00
Architectural Iron and Steel.	8vo,	3 50
Compound Riveted Girders as Applied in Buildings.	8vo,	2 00
Planning and Construction of High Office Buildings.	8vo,	3 50
Skeleton Construction in Buildings.	8vo,	3 00
Brigg's Modern American School Buildings.	8vo,	4 00
Carpenter's Heating and Ventilating of Buildings.	8vo,	4 00
Freitag's Architectural Engineering.	8vo,	3 50
Fireproofing of Steel Buildings.	8vo,	2 50
French and Ives's Stereotomy.	8vo,	2 50

Gerhard's Guide to Sanitary House-inspection.....	16mo,	1 00
Theatre Fires and Panics.....	12mo,	1 50
*Greene's Structural Mechanics.....	8vo,	2 50
Holly's Carpenters' and Joiners' Handbook.....	18mo,	75
Johnson's Statics by Algebraic and Graphic Methods.....	8vo,	2 00
Kidder's Architects' and Builders' Pocket-book. Rewritten Edition. 16mo,mor.,	5 00	
Merrill's Stones for Building and Decoration.....	8vo,	5 00
Non-metallic Minerals: Their Occurrence and Uses.....	8vo,	4 00
Monckton's Stair-building.....	4to,	4 00
Patton's Practical Treatise on Foundations.....	8vo,	5 00
Peabody's Naval Architecture.....	8vo,	7 50
Richey's Handbook for Superintendents of Construction.....	16mo, mor.,	4 00
Sabin's Industrial and Artistic Technology of Paints and Varnish.....	8vo,	3 00
Siebert and Biggin's Modern Stone-cutting and Masonry.....	8vo,	1 50
Snow's Principal Species of Wood.....	8vo,	3 50
Sondericker's Graphic Statics with Applications to Trusses, Beams, and Arches.	8vo,	2 00
Towne's Locks and Builders' Hardware.....	18mo, morocco,	3 00
Wait's Engineering and Architectural Jurisprudence.....	8vo,	6 00
	Sheep,	6 50
Law of Operations Preliminary to Construction in Engineering and Archi- tecture.....	8vo,	5 00
	Sheep,	5 50
Law of Contracts.....	8vo,	3 00
Wood's Rustless Coatings: Corrosion and Electrolysis of Iron and Steel. 8vo,	4 00	
Worcester and Atkinson's Small Hospitals, Establishment and Maintenance, Suggestions for Hospital Architecture, with Plans for a Small Hospital.	12mo,	1 25
The World's Columbian Exposition of 1893.....	Large 4to,	1 00

ARMY AND NAVY.

Bernadou's Smokeless Powder, Nitro-cellulose, and the Theory of the Cellulose Molecule.....	12mo,	2 50
*Bruff's Text-book Ordnance and Gunnery.....	8vo,	6 00
Chase's Screw Propellers and Marine Propulsion.....	8vo,	3 00
Cloke's Gunner's Examiner.....	8vo,	1 50
Craig's Azimuth.....	4to,	3 50
Crehore and Squier's Polarizing Photo-chronograph.....	8vo,	3 00
*Davis's Elements of Law.....	8vo,	2 50
* Treatise on the Military Law of United States.....	8vo,	7 00
	Sheep,	7 50
De Brack's Cavalry Outposts Duties. (Carr.).....	24mo, morocco,	2 00
Dietz's Soldier's First Aid Handbook.....	16mo, morocco,	1 25
*Dredge's Modern French Artillery.....	4to, half morocco,	15 00
Durand's Resistance and Propulsion of Ships.....	8vo,	5 00
*Dyer's Handbook of Light Artillery.....	12mo,	3 00
Eissler's Modern High Explosives.....	8vo,	4 00
*Fiebigger's Text-book on Field Fortification.....	Small 8vo,	2 00
Hamilton's The Gunner's Catechism.....	18mo,	1 00
*Hoff's Elementary Naval Tactics.....	8vo,	1 50
Ingalls's Handbook of Problems in Direct Fire.....	8vo,	4 00
* Ballistic Tables.....	8vo,	1 50
*Lyons's Treatise on Electromagnetic Phenomena. Vols. I. and II. 8vo, each,	6 00	
*Mahan's Permanent Fortifications. (Mercur.).....	8vo, half morocco,	7 50
Manual for Courts-martial.....	16mo, morocco,	1 50
*Mercur's Attack of Fortified Places.....	12mo,	2 00
* Elements of the Art of War.....	8vo,	4 00

Metcalf's Cost of Manufactures—And the Administration of Workshops.	8vo,	5 00
* Ordnance and Gunnery. 2 vols.	12mo,	5 00
Murray's Infantry Drill Regulations.	18mo, paper,	10
Nixon's Adjutants' Manual.	24mo,	1 00
Peabody's Naval Architecture.	8vo,	7 50
* Phelps's Practical Marine Surveying.	8vo,	2 50
Powell's Army Officer's Examiner.	12mo,	4 00
Sharpe's Art of Subsisting Armies in War.	18mo, morocco,	1 50
* Walke's Lectures on Explosives.	8vo,	4 00
* Wheeler's Siege Operations and Military Mining.	8vo,	2 00
Winthrop's Abridgment of Military Law.	12mo,	2 50
Woodhull's Notes on Military Hygiene.	16mo,	1 50
Young's Simple Elements of Navigation.	16mo, morocco,	2 00

ASSAYING.

Fletcher's Practical Instructions in Quantitative Assaying with the Blowpipe.	12mo, morocco,	1 50
Furman's Manual of Practical Assaying.	8vo,	3 00
Lodge's Notes on Assaying and Metallurgical Laboratory Experiments.	8vo,	3 00
Low's Technical Methods of Ore Analysis.	8vo,	3 00
Miller's Manual of Assaying.	12mo,	1 00
Minet's Production of Aluminum and its Industrial Use. (Waldo.).	12mo,	2 50
O'Driscoll's Notes on the Treatment of Gold Ores.	8vo,	2 00
Ricketts and Miller's Notes on Assaying.	8vo,	3 00
Robine and Lenglen's Cyanide Industry. (Le Clerc.).	8vo,	
Ulke's Modern Electrolytic Copper Refining.	8vo,	3 00
Wilson's Cyanide Processes.	12mo,	1 50
Chlorination Process.	12mo,	1 50

ASTRONOMY.

Comstock's Field Astronomy for Engineers.	8vo,	2 50
Craig's Azimuth.	4to,	3 50
Doolittle's Treatise on Practical Astronomy.	8vo,	4 00
Gore's Elements of Geodesy.	8vo,	2 50
Hayford's Text-book of Geodetic Astronomy.	8vo,	3 00
Merriman's Elements of Precise Surveying and Geodesy.	8vo,	2 50
* Michie and Harlow's Practical Astronomy.	8vo,	3 00
* White's Elements of Theoretical and Descriptive Astronomy	12mo,	2 00

BOTANY.

Davenport's Statistical Methods, with Special Reference to Biological Variation.	16mo, morocco,	1 25
Thomé and Bennett's Structural and Physiological Botany.	16mo,	2 25
Westermaier's Compendium of General Botany. (Schneider.).	8vo,	2 00

CHEMISTRY.

Adrianse's Laboratory Calculations and Specific Gravity Tables.	12mo,	1 25
Allen's Tables for Iron Analysis.	8vo,	3 00
Arnold's Compendium of Chemistry. (Mandel.).	Small 8vo,	3 50
Austen's Notes for Chemical Students	12mo,	1 50
Bernadou's Smokeless Powder.—Nitro-cellulose, and Theory of the Cellulose Molecule.	12mo,	2 50
* Browning's Introduction to the Rarer Elements.	8vo,	1 50

Brush and Penfield's Manual of Determinative Mineralogy.	8vo,	4 00
Classen's Quantitative Chemical Analysis by Electrolysis. (Boltwood.)...	8vo,	3 00
Cohn's Indicators and Test-papers.	12mo,	2 00
Tests and Reagents.	8vo,	3 00
Crafts's Short Course in Qualitative Chemical Analysis. (Schaeffer.)...	12mo,	1 50
Dolezalek's Theory of the Lead Accumulator (Storage Battery). (Von Ende.)...	12mo,	2 50
Drechsel's Chemical Reactions. (Merrill.)...	12mo,	1 25
Duhem's Thermodynamics and Chemistry. (Burgess.)...	8vo,	4 00
Eissler's Modern High Explosives.	8vo,	4 00
Effront's Enzymes and their Applications. (Prescott.)...	8vo,	3 00
Erdmann's Introduction to Chemical Preparations. (Dunlap.)...	12mo,	1 25
Fletcher's Practical Instructions in Quantitative Assaying with the Blowpipe.	12mo, morocco,	1 50
Fowler's Sewage Works Analyses.	12mo,	2 00
Fresenius's Manual of Qualitative Chemical Analysis. (Wells.)...	8vo,	5 00
Manual of Qualitative Chemical Analysis, Part I, Descriptive. (Wells.)	8vo,	3 00
System of Instruction in Quantitative Chemical Analysis. (Cohn.)	2 vols.	8vo, 12 50
Fuertes's Water and Public Health.	12mo,	1 50
Furman's Manual of Practical Assaying.	8vo,	3 00
* Getman's Exercises in Physical Chemistry.	12mo,	2 00
Gill's Gas and Fuel Analysis for Engineers.	12mo,	1 25
Grotenfelt's Principles of Modern Dairy Practice. (Woll.)...	12mo,	2 00
Hammarsten's Text-book of Physiological Chemistry. (Mandel.)...	8vo,	4 00
Helm's Principles of Mathematical Chemistry. (Morgan.)...	12mo,	1 50
Hering's Ready Reference Tables (Conversion Factors).	16mo, morocco,	2 50
Hind's Inorganic Chemistry.	8vo,	3 00
* Laboratory Manual for Students.	12mo,	1 00
Holleman's Text-book of Inorganic Chemistry. (Cooper.)...	8vo,	2 50
Text-book of Organic Chemistry. (Walker and Mott.)...	8vo,	2 50
* Laboratory Manual of Organic Chemistry. (Walker.)...	12mo,	1 00
Hopkins's Oil-chemists' Handbook.	8vo,	3 00
Jackson's Directions for Laboratory Work in Physiological Chemistry.	8vo,	1 25
Keep's Cast Iron.	8vo,	2 50
Ladd's Manual of Quantitative Chemical Analysis.	12mo,	1 00
Landauer's Spectrum Analysis. (Tingle.)...	8vo,	3 00
* Langworthy and Austen. The Occurrence of Aluminium in Vegetable Products, Animal Products, and Natural Waters.	8vo,	2 00
Lassar-Cohn's Practical Urinary Analysis. (Lorenz.)...	12mo,	1 00
Application of Some General Reactions to Investigations in Organic Chemistry. (Tingle.)...	12mo,	1 00
Leach's The Inspection and Analysis of Food with Special Reference to State Control.	8vo,	7 50
Löb's Electrochemistry of Organic Compounds. (Lorenz.)...	8vo,	3 00
Lodge's Notes on Assaying and Metallurgical Laboratory Experiments.	8vo,	3 00
Low's Technical Method of Ore Analysis.	8vo,	3 00
Lunge's Techno-chemical Analysis. (Cohn.)...	12mo,	1 00
Mandel's Handbook for Bio-chemical Laboratory.	12mo,	1 50
* Martin's Laboratory Guide to Qualitative Analysis with the Blowpipe.	12mo,	60
Mason's Water-supply. (Considered Principally from a Sanitary Standpoint.)	3d Edition, Rewritten.	8vo, 4 00
Examination of Water. (Chemical and Bacteriological.)...	12mo,	1 25
Matthew's The Textile Fibres.	8vo,	3 50
Meyer's Determination of Radicles in Carbon Compounds. (Tingle.)...	12mo,	1 00
Miller's Manual of Assaying.	12mo,	1 00
Minet's Production of Aluminum and its Industrial Use. (Waldo.)...	12mo,	2 50
Mixter's Elementary Text-book of Chemistry.	12mo,	1 50
Morgan's Elements of Physical Chemistry.	12mo,	3 00
* Physical Chemistry for Electrical Engineers.	12mo,	1 50

CIVIL ENGINEERING.

BRIDGES AND ROOFS. HYDRAULICS. MATERIALS OF ENGINEERING. RAILWAY ENGINEERING.

Baker's Engineers' Surveying Instruments.	12mo,	3 00
Bixby's Graphical Computing Table.	Paper 19½ × 24½ inches.	25
** Burr's Ancient and Modern Engineering and the Isthmian Canal. (Postage, 27 cents additional.)		
Comstock's Field Astronomy for Engineers.	8vo,	3 50
Davis's Elevation and Stadia Tables.	8vo,	2 50
Elliott's Engineering for Land Drainage.	8vo,	1 00
Practical Farm Drainage.	12mo,	1 50
* Fiebegeer's Treatise on Civil Engineering.	12mo,	1 00
Folwell's Sewerage. (Designing and Maintenance.)	8vo,	5 00
Freitag's Architectural Engineering. 2d Edition, Rewritten	8vo,	3 00
French and Ives's Stereotomy.	8vo,	3 50
Goodhue's Municipal Improvements.	8vo,	2 50
Goodrich's Economic Disposal of Towns' Refuse.	12mo,	1 75
Gore's Elements of Geodesy.	8vo,	3 50
Hayford's Text-book of Geodetic Astronomy.	8vo,	2 50
Hering's Ready Reference Tables (Conversion Factors).	8vo,	3 00
Howe's Retaining Walls for Earth.	16mo, morocco,	2 50
Johnson's (J. B.) Theory and Practice of Surveying.	12mo,	1 25
Johnson's (L. J.) Statics by Algebraic and Graphic Methods.	Small 8vo,	4 00
Laplace's Philosophical Essay on Probabilities. (Truscott and Emory.)	8vo,	2 00
Mahan's Treatise on Civil Engineering. (1873.) (Wood.)	8vo,	2 00
* Descriptive Geometry.	8vo,	5 00
Merriman's Elements of Precise Surveying and Geodesy.	8vo,	1 50
Merriman and Brooks's Handbook for Surveyors.	8vo,	2 50
Nugent's Plane Surveying.	16mo, morocco,	7 00
Ogden's Sewer Design.	8vo,	3 50
Patton's Treatise on Civil Engineering.	12mo,	2 00
Reed's Topographical Drawing and Sketching	8vo half leather,	7 50
Rideal's Sewage and the Bacterial Purification of Sewage.	4to,	5 00
Siebert and Biggin's Modern Stone-cutting and Masonry.	8vo,	3 50
Smith's Manual of Topographical Drawing. (McMillan.)	8vo,	1 50
Sondericker's Graphic Statics, with Applications to Trusses, Beams, and Arches.	8vo,	2 50
Taylor and Thompson's Treatise on Concrete, Plain and Reinforced.	8vo,	2 00
* Trautwine's Civil Engineer's Pocket-book.	8vo,	5 00
Wait's Engineering and Architectural Jurisprudence.	16mo, morocco,	6 00
Law of Operations Preliminary to Construction in Engineering and Architecture.	8vo,	6 50
Law of Contracts.	8vo,	5 00
Warren's Stereotomy—Problems in Stone-cutting.	8vo,	3 00
Webb's Problems in the Use and Adjustment of Engineering Instruments.	8vo,	2 50
Wilson's Topographic Surveying.	16mo, morocco,	1 25
	8vo,	3 50

BRIDGES AND ROOFS.

Boller's Practical Treatise on the Construction of Iron Highway Bridges.	8vo,	2 00
* Thames River Bridge.	4to, paper,	5 00
Burr's Course on the Stresses in Bridges and Roof Trusses, Arched Ribs, and Suspension Bridges.	8vo,	3 50

Burr and Falk's Influence Lines for Bridge and Roof Computations.	8vo,	3 00
Design and Construction of Metallic Bridges.	8vo,	5 00
Du Bois's Mechanics of Engineering. Vol. II.	Small 4to,	10 00
Foster's Treatise on Wooden Trestle Bridges.	4to,	5 00
Fowler's Ordinary Foundations.	8vo,	3 50
Greene's Roof Trusses.	8vo,	1 25
Bridge Trusses.	8vo,	2 50
Arches in Wood, Iron, and Stone.	8vo,	2 50
Howe's Treatise on Arches.	8vo,	4 00
Design of Simple Roof-trusses in Wood and Steel.	8vo,	2 00
Johnson, Bryan, and Turneure's Theory and Practice in the Designing of Modern Framed Structures.	Small 4to,	10 00
Merriman and Jacoby's Text-book on Roofs and Bridges:		
Part I. Stresses in Simple Trusses.	8vo,	2 50
Part II. Graphic Statics.	8vo,	2 50
Part III. Bridge Design.	8vo,	2 50
Part IV. Higher Structures.	8vo,	2 50
Morison's Memphis Bridge.	4to,	10 00
Waddell's De Pontibus, a Pocket-book for Bridge Engineers. . 16mo, morocco,		2 00
Specifications for Steel Bridges.	12mo,	1 25
Wright's Designing of Draw-spans. Two parts in one volume.	8vo,	3 50

HYDRAULICS.

Bazin's Experiments upon the Contraction of the Liquid Vein Issuing from an Orifice. (Trautwine.)	8vo,	2 00
Bovey's Treatise on Hydraulics.	8vo,	5 00
Church's Mechanics of Engineering.	8vo,	6 00
Diagrams of Mean Velocity of Water in Open Channels.	paper,	1 50
Hydraulic Motors.	8vo,	2 00
Coffin's Graphical Solution of Hydraulic Problems.	16mo, morocco,	2 50
Fletcher's Dynamometers, and the Measurement of Power.	12mo,	3 00
Folwell's Water-supply Engineering.	8vo,	4 00
Frizell's Water-power.	8vo,	5 00
Fuertes's Water and Public Health.	12mo,	1 50
Water-filtration Works.	12mo,	2 50
Ganguillet and Kutter's General Formula for the Uniform Flow of Water in Rivers and Other Channels. (Hering and Trautwine.)	8vo,	4 00
Hazen's Filtration of Public Water-supply.	8vo,	3 00
Hazlehurst's Towers and Tanks for Water-works.	8vo,	2 50
Herschel's 115 Experiments on the Carrying Capacity of Large, Riveted, Metal Conduits.	8vo,	2 00
Mason's Water-supply. (Considered Principally from a Sanitary Standpoint.)	8vo,	4 00
Merriman's Treatise on Hydraulics.	8vo,	5 00
* Michie's Elements of Analytical Mechanics.	8vo,	4 00
Schuyler's Reservoirs for Irrigation, Water-power, and Domestic Water- supply.	Large 8vo,	5 00
** Thomas and Watt's Improvement of Rivers. (Post., 44c. additional.) . 4to,		6 00
Turneure and Russell's Public Water-supplies.	8vo,	5 00
Wegmann's Design and Construction of Dams.	4to,	5 00
Water-supply of the City of New York from 1658 to 1895.	4to,	10 00
Williams and Hazen's Hydraulic Tables.	8vo,	1 50
Wilson's Irrigation Engineering.	Small 8vo,	4 00
Wolff's Windmill as a Prime Mover.	8vo,	3 00
Wood's Turbines.	8vo,	2 50
Elements of Analytical Mechanics.	8vo,	3 00

MATERIALS OF ENGINEERING.

Baker's Treatise on Masonry Construction.	8vo,	5 00
Roads and Pavements.	8vo,	5 00
Black's United States Public Works.	Oblong 4to,	5 00
* Bovey's Strength of Materials and Theory of Structures.	8vo,	7 50
Burr's Elasticity and Resistance of the Materials of Engineering.	8vo,	7 50
Byrne's Highway Construction.	8vo,	5 00
Inspection of the Materials and Workmanship Employed in Construction.	16mo,	3 00
Church's Mechanics of Engineering.	8vo,	6 00
Du Bois's Mechanics of Engineering. Vol. I.	Small 4to,	7 50
*Eckel's Cements, Limes, and Plasters.	8vo,	6 00
Johnson's Materials of Construction.	Large 8vo,	6 00
Fowler's Ordinary Foundations.	8vo,	3 50
* Greene's Structural Mechanics.	8vo,	2 50
Keep's Cast Iron.	8vo,	2 50
Lanza's Applied Mechanics.	8vo,	7 50
Marten's Handbook on Testing Materials. (Henning.) 2 vols.	8vo,	7 50
Maurer's Technical Mechanics.	8vo,	4 00
Merrill's Stones for Building and Decoration.	8vo,	5 00
Merriman's Mechanics of Materials.	8vo,	5 00
Strength of Materials.	12mo,	1 00
Metcalfe's Steel. A Manual for Steel-users.	12mo,	2 00
Patton's Practical Treatise on Foundations.	8vo,	5 00
Richardson's Modern Asphalt Pavements.	8vo,	3 00
Richey's Handbook for Superintendents of Construction.	16mo, mor.,	4 00
Rockwell's Roads and Pavements in France.	12mo,	1 25
Sabin's Industrial and Artistic Technology of Paints and Varnish.	8vo,	3 00
Smith's Materials of Machines.	12mo,	1 00
Snow's Principal Species of Wood.	8vo,	3 50
Spalding's Hydraulic Cement.	12mo,	2 00
Text-book on Roads and Pavements.	12mo,	2 00
Taylor and Thompson's Treatise on Concrete, Plain and Reinforced.	8vo,	5 00
Thurston's Materials of Engineering. 3 Parts.	8vo,	8 00
Part I. Non-metallic Materials of Engineering and Metallurgy.	8vo,	2 00
Part II. Iron and Steel.	8vo,	3 50
Part III. A Treatise on Brasses, Bronzes, and Other Alloys and their Constituents.	8vo,	2 50
Thurston's Text-book of the Materials of Construction.	8vo,	5 00
Tillson's Street Pavements and Paving Materials.	8vo,	4 00
Waddell's De Pontibus. (A Pocket-book for Bridge Engineers.) .	16mo, mor.,	2 00
Specifications for Steel Bridges.	12mo,	1 25
Wood's (De V.) Treatise on the Resistance of Materials, and an Appendix on the Preservation of Timber.	8vo,	2 00
Wood's (De V.) Elements of Analytical Mechanics.	8vo,	3 00
Wood's (M. P.) Rustless Coatings: Corrosion and Electrolysis of Iron and Steel.	8vo,	4 00

. RAILWAY ENGINEERING.

Andrew's Handbook for Street Railway Engineers.	3x5 inches, morocco,	1 25
Berg's Buildings and Structures of American Railroads.	4to,	5 00
Brook's Handbook of Street Railroad Location.	16mo, morocco,	1 50
Butt's Civil Engineer's Field-book.	16mo, morocco,	2 50
Crandall's Transition Curve.	16mo, morocco,	1 50
Railway and Other Earthwork Tables.	8vo,	1 50
Dawson's "Engineering" and Electric Traction Pocket-book.	16mo, morocco,	5 00

Dredge's History of the Pennsylvania Railroad: (1879).....	Paper,	5 00
* Drinker's Tunnelling, Explosive Compounds, and Rock Drills.....	4to, half mor.,	25 00
Fisher's Table of Cubic Yards.....	Cardboard,	25
Godwin's Railroad Engineers' Field-book and Explorers' Guide.....	16mo, mor.,	2 50
Howard's Transition Curve Field-book.....	16mo, morocco,	1 50
Hudson's Tables for Calculating the Cubic Contents of Excavations and Embankments.....	8vo,	1 00
Molitor and Beard's Manual for Resident Engineers.....	16mo,	1 00
Nagle's Field Manual for Railroad Engineers.....	16mo, morocco,	3 00
Philbrick's Field Manual for Engineers.....	16mo, morocco,	3 00
Searles's Field Engineering.....	16mo, morocco,	3 00
Railroad Spiral.....	16mo, morocco,	1 50
Taylor's Prismoidal Formulæ and Earthwork.....	8vo,	1 50
* Trautwine's Method of Calculating the Cube Contents of Excavations and Embankments by the Aid of Diagrams.....	8vo,	2 00
The Field Practice of Laying Out Circular Curves for Railroads.....	12mo, morocco,	2 50
Cross-section Sheet.....	Paper,	25
Webb's Railroad Construction.....	16mo, morocco,	5 00
Wellington's Economic Theory of the Location of Railways.....	Small 8vo,	5 00

DRAWING.

Barr's Kinematics of Machinery.....	8vo,	2 50
* Bartlett's Mechanical Drawing.....	8vo,	3 00
* " " " " Abridged Ed.....	8vo,	1 50
Coolidge's Manual of Drawing.....	8vo, paper	1 00
Coolidge and Freeman's Elements of General Drafting for Mechanical Engineers.....	Oblong 4to,	2 50
Durley's Kinematics of Machines.....	8vo,	4 00
Emch's Introduction to Projective Geometry and its Applications.....	8vo,	2 50
Hill's Text-book on Shades and Shadows, and Perspective.....	8vo,	2 00
Jamison's Elements of Mechanical Drawing.....	8vo,	2 50
Advanced Mechanical Drawing.....	8vo,	2 00
Jones's Machine Design:		
Part I. Kinematics of Machinery.....	8vo,	1 50
Part II. Form, Strength, and Proportions of Parts.....	8vo,	3 00
MacCord's Elements of Descriptive Geometry.....	8vo,	3 00
Kinematics; or, Practical Mechanism.....	8vo,	5 00
Mechanical Drawing.....	4to,	4 00
Velocity Diagrams.....	8vo,	1 50
MacLeod's Descriptive Geometry.....	Small 8vo,	1 50
* Mahan's Descriptive Geometry and Stone-cutting.....	8vo,	1 50
Industrial Drawing. (Thompson.).....	8vo,	3 50
Moyer's Descriptive Geometry.....	8vo,	2 00
Reed's Topographical Drawing and Sketching.....	4to,	5 00
Reid's Course in Mechanical Drawing.....	8vo,	2 00
Text-book of Mechanical Drawing and Elementary Machine Design.....	8vo,	3 00
Robinson's Principles of Mechanism.....	8vo,	3 00
Schwamb and Merrill's Elements of Mechanism.....	8vo,	3 00
Smith's (R. S.) Manual of Topographical Drawing. (McMillan.).....	8vo,	2 50
Smith (A. W.) and Marx's Machine Design.....	8vo,	3 00
Warren's Elements of Plane and Solid Free-hand Geometrical Drawing.....	12mo,	1 00
Drafting Instruments and Operations.....	12mo,	1 25
Manual of Elementary Projection Drawing.....	12mo,	1 50
Manual of Elementary Problems in the Linear Perspective of Form and Shadow.....	12mo,	1 00
Plane Problems in Elementary Geometry.....	12mo,	1 25

Warren's Primary Geometry.....	12mo,	75
Elements of Descriptive Geometry, Shadows, and Perspective.....	8vo,	3 50
General Problems of Shades and Shadows.....	8vo,	3 00
Elements of Machine Construction and Drawing.....	8vo,	7 50
Problems, Theorems, and Examples in Descriptive Geometry.....	8vo,	2 50
Weisbach's Kinematics and Power of Transmission. (Hermann and Klein.).....	8vo,	5 00
Whelpley's Practical Instruction in the Art of Letter Engraving.....	12mo,	2 00
Wilson's (H. M.) Topographic Surveying.....	8vo,	3 50
Wilson's (V. T.) Free-hand Perspective.....	8vo,	2 50
Wilson's (V. T.) Free-hand Lettering.....	8vo,	1 00
Woolf's Elementary Course in Descriptive Geometry.....	Large 8vo,	3 00

ELECTRICITY AND PHYSICS.

Anthony and Brackett's Text-book of Physics. (Magie.).....	Small 8vo,	3 00
Anthony's Lecture-notes on the Theory of Electrical Measurements.....	12mo,	1 00
Benjamin's History of Electricity.....	8vo,	3 00
Voltaic Cell.....	8vo,	3 00
Classen's Quantitative Chemical Analysis by Electrolysis. (Boltwood.).....	8vo,	3 00
Crehore and Squier's Polarizing Photo-chronograph.....	8vo,	3 00
Dawson's "Engineering" and Electric Traction Pocket-book.....	16mo, morocco,	5 00
Dolezalek's Theory of the Lead Accumulator (Storage Battery). (Von Ende.).....	12mo,	2 50
Duhem's Thermodynamics and Chemistry. (Burgess.).....	8vo,	4 00
Flather's Dynamometers, and the Measurement of Power.....	12mo,	3 00
Gilbert's De Magnete. (Mottelay.).....	8vo,	2 50
Hanchett's Alternating Currents Explained.....	12mo,	1 00
Hering's Ready Reference Tables (Conversion Factors).....	16mo, morocco,	2 50
Holman's Precision of Measurements.....	8vo,	2 00
Telescopic Mirror-scale Method, Adjustments, and Tests.....	Large 8vo,	75
Kinzbrunner's Testing of Continuous-current Machines.....	8vo,	2 00
Landauer's Spectrum Analysis. (Tingle.).....	8vo,	3 00
Le Chatelier's High-temperature Measurements. (Boudouard—Burgess.).....	12mo,	3 00
Lob's Electrochemistry of Organic Compounds. (Lorenz.).....	8vo,	3 00
* Lyons's Treatise on Electromagnetic Phenomena. Vols. I. and II. 8vo, each.....	8vo,	6 00
* Michie's Elements of Wave Motion Relating to Sound and Light.....	8vo,	4 00
Niaudet's Elementary Treatise on Electric Batteries. (Fishback.).....	12mo,	2 50
* Rosenberg's Electrical Engineering. (Haldane Gee—Kinzbrunner.).....	8vo,	1 50
Ryan, Norris, and Hoxie's Electrical Machinery. Vol. I.....	8vo,	2 50
Thurston's Stationary Steam-engines.....	8vo,	2 50
* Tillman's Elementary Lessons in Heat.....	8vo,	1 50
Tory and Pitcher's Manual of Laboratory Physics.....	Small 8vo,	2 00
Ulke's Modern Electrolytic Copper Refining.....	8vo,	3 00

LAW.

* Davis's Elements of Law.....	8vo,	2 50
* Treatise on the Military Law of United States.....	8vo,	7 00
*	Sheep,	7 50
Manual for Courts-martial.....	16mo, morocco,	1 50
Wait's Engineering and Architectural Jurisprudence.....	8vo,	6 00
.....	Sheep,	6 50
Law of Operations Preliminary to Construction in Engineering and Architecture.....	8vo	5 00
.....	Sheep,	5 50
Law of Contracts.....	8vo,	3 00
Winthrop's Abridgment of Military Law.....	12mo.	2 50

MANUFACTURES.

Bernadou's Smokeless Powder—Nitro-cellulose and Theory of the Cellulose Molecule.....	12mo,	2 50
Bolland's Iron Founder.....	12mo,	2 50
"The Iron Founder," Supplement.....	12mo,	2 50
Encyclopedia of Founding and Dictionary of Foundry Terms Used in the Practice of Moulding.....	12mo,	3 00
Eissler's Modern High Explosives.....	8vo,	4 00
Effront's Enzymes and their Applications. (Prescott.).....	8vo,	3 00
Fitzgerald's Boston Machinist.....	12mo,	1 00
Ford's Boiler Making for Boiler Makers.....	18mo,	1 00
Hopkin's Oil-chemists' Handbook.....	8vo,	3 00
Keep's Cast Iron.....	8vo,	2 50
Leach's The Inspection and Analysis of Food with Special Reference to State Control.....	Large 8vo,	7 50
Matthews's The Textile Fibres.....	8vo,	3 50
Metcalf's Steel. A Manual for Steel-users.....	12mo,	2 00
Metcalfe's Cost of Manufactures—And the Administration of Workshops.....	8vo,	5 00
Meyer's Modern Locomotive Construction.....	4to,	10 00
Morse's Calculations used in Cane-sugar Factories.....	16mo, morocco,	1 50
* Reising's Guide to Piece-dyeing.....	8vo,	25 00
Sabin's Industrial and Artistic Technology of Paints and Varnish.....	8vo,	3 00
Smith's Press-working of Metals.....	8vo,	3 00
Spalding's Hydraulic Cement.....	12mo,	2 00
Spencer's Handbook for Chemists of Beet-sugar Houses.....	16mo, morocco,	3 00
Handbook for Cane Sugar Manufacturers.....	16mo, morocco,	3 00
Taylor and Thompson's Treatise on Concrete, Plain and Reinforced.....	8vo,	5 00
Thurston's Manual of Steam-boilers, their Designs, Construction and Operation.....	8vo,	5 00
* Walke's Lectures on Explosives.....	8vo,	4 00
Ware's Beet-sugar Manufacture and Refining.....	Small 8vo,	4 00
West's American Foundry Practice.....	12mo,	2 50
Moulder's Text-book.....	12mo,	2 50
Wolff's Windmill as a Prime Mover.....	8vo,	3 00
Wood's Rustless Coatings: Corrosion and Electrolysis of Iron and Steel.....	8vo,	4 00

MATHEMATICS.

Baker's Elliptic Functions.....	8vo,	1 50
* Bass's Elements of Differential Calculus.....	12mo,	4 00
Briggs's Elements of Plane Analytic Geometry.....	12mo,	1 00
Compton's Manual of Logarithmic Computations.....	12mo,	1 50
Davis's Introduction to the Logic of Algebra.....	8vo,	1 50
* Dickson's College Algebra.....	Large 12mo,	1 50
* Introduction to the Theory of Algebraic Equations.....	Large 12mo,	1 25
Emch's Introduction to Projective Geometry and its Applications.....	8vo,	2 50
Halsted's Elements of Geometry.....	8vo,	1 75
Elementary Synthetic Geometry.....	8vo,	1 50
Rational Geometry.....	12mo,	1 75
* Johnson's (J. B.) Three-place Logarithmic Tables: Vest-pocket size, paper.....		15
100 copies for.....		5 00
* Mounted on heavy cardboard, 8×10 inches.....		25
10 copies for.....		2 00
Johnson's (W. W.) Elementary Treatise on Differential Calculus.....	Small 8vo,	3 00
Johnson's (W. W.) Elementary Treatise on the Integral Calculus.....	Small 8vo,	1 50

Johnson's (W. W.) Curve Tracing in Cartesian Co-ordinates.....	12mo,	1 00
Johnson's (W. W.) Treatise on Ordinary and Partial Differential Equations.		
	Small 8vo,	3 50
Johnson's (W. W.) Theory of Errors and the Method of Least Squares.....	12mo,	1 50
* Johnson's (W. W.) Theoretical Mechanics.....	12mo,	3 00
Laplace's Philosophical Essay on Probabilities. (Truscott and Emory.).....	12mo,	2 00
* Ludlow and Bass. Elements of Trigonometry and Logarithmic and Other Tables.....	8vo,	3 00
Trigonometry and Tables published separately.....	Each,	2 00
* Ludlow's Logarithmic and Trigonometric Tables.....	8vo,	1 00
Mathematical Monographs. Edited by Mansfield Merriman and Robert S. Woodward.....	Octavo, each	1 00
No. 1. History of Modern Mathematics, by David Eugene Smith.		
No. 2. Synthetic Projective Geometry, by George Bruce Halsted.		
No. 3. Determinants, by Laenas Gifford Weld. No. 4. Hyper- bolic Functions, by James McMahon. No. 5. Harmonic Func- tions, by William E. Byerly. No. 6. Grassmann's Space Analysis, by Edward W. Hyde. No. 7. Probability and Theory of Errors, by Robert S. Woodward. No. 8. Vector Analysis and Quaternions, by Alexander Macfarlane. No. 9. Differential Equations, by William Woolsey Johnson. No. 10. The Solution of Equations, by Mansfield Merriman. No. 11. Functions of a Complex Variable, by Thomas S. Fiske.		
Maurer's Technical Mechanics.....	8vo,	4 00
Merriman and Woodward's Higher Mathematics.....	8vo,	5 00
Merriman's Method of Least Squares.....	8vo,	2 00
Rice and Johnson's Elementary Treatise on the Differential Calculus.....	Sm. 8vo,	3 00
Differential and Integral Calculus. 2 vols. in one.....	Small 8vo,	2 50
Wood's Elements of Co-ordinate Geometry.....	8vo,	2 00
Trigonometry: Analytical, Plane, and Spherical.....	12mo,	1 00

MECHANICAL ENGINEERING.

MATERIALS OF ENGINEERING, STEAM-ENGINES AND BOILERS.

Bacon's Forge Practice.....	12mo,	1 50
Baldwin's Steam Heating for Buildings.....	12mo,	2 50
Barr's Kinematics of Machinery.....	8vo,	2 50
* Bartlett's Mechanical Drawing.....	8vo,	3 00
* " " Abridged Ed.....	8vo,	1 50
Benjamin's Wrinkles and Recipes.....	12mo,	2 00
Carpenter's Experimental Engineering.....	8vo,	6 00
Heating and Ventilating Buildings.....	8vo,	4 00
Cary's Smoke Suppression in Plants using Bituminous Coal. (In Prepara- tion.)		
Clerk's Gas and Oil Engine.....	Small 8vo,	4 00
Coolidge's Manual of Drawing.....	8vo, paper,	1 00
Coolidge and Freeman's Elements of General Drafting for Mechanical En- gineers.....	Oblong 4to,	2 50
Cromwell's Treatise on Toothed Gearing.....	12mo,	1 50
Treatise on Belts and Pulleys.....	12mo,	1 50
Durley's Kinematics of Machines.....	8vo,	4 00
Flather's Dynamometers and the Measurement of Power.....	12mo,	3 00
Rope Driving.....	12mo,	2 00
Gill's Gas and Fuel Analysis for Engineers.....	12mo,	1 25
Hall's Car Lubrication.....	12mo,	1 00
Hering's Ready Reference Tables (Conversion Factors).....	16mo, morocco,	2 50

Hutton's The Gas Engine.	8vo,	5 00
Jamison's Mechanical Drawing.	8vo,	2 50
Jones's Machine Design:		
Part I. Kinematics of Machinery.	8vo,	1 50
Part II. Form, Strength, and Proportions of Parts.	8vo,	3 00
Kent's Mechanical Engineers' Pocket-book.	16mo, morocco,	5 00
Kerr's Power and Power Transmission.	8vo,	2 00
Leonard's Machine Shop, Tools, and Methods.	8vo,	4 00
* Lorenz's Modern Refrigerating Machinery. (Pope, Haven, and Dean.)	8vo,	4 00
MacCord's Kinematics; or, Practical Mechanism.	8vo,	5 00
Mechanical Drawing.	4to,	4 00
Velocity Diagrams.	8vo,	1 50
MacFarland's Standard Reduction Factors for Gases.	8vo,	1 50
Mahan's Industrial Drawing. (Thompson.)	8vo,	3 50
Poole's Calorific Power of Fuels.	8vo,	3 00
Reid's Course in Mechanical Drawing.	8vo,	2 00
Text-book of Mechanical Drawing and Elementary Machine Design.	8vo,	3 00
Richard's Compressed Air	12mo,	1 50
Robinson's Principles of Mechanism.	8vo,	3 00
Schwamb and Merrill's Elements of Mechanism.	8vo,	3 00
Smith's (O.) Press-working of Metals.	8vo,	3 00
Smith (A. W.) and Marx's Machine Design.	8vo,	3 00
Thurston's Treatise on Friction and Lost Work in Machinery and Mill Work.	8vo,	3 00
Animal as a Machine and Prime Motor, and the Laws of Energetics.	12mo,	1 00
Warren's Elements of Machine Construction and Drawing.	8vo,	7 50
Weisbach's Kinematics and the Power of Transmission. (Herrmann—Klein.)	8vo,	5 00
Machinery of Transmission and Governors. (Herrmann—Klein.)	8vo,	5 00
Wolff's Windmill as a Prime Mover.	8vo,	3 00
Wood's Turbines.	8vo,	2 50

MATERIALS OF ENGINEERING.

* Bovey's Strength of Materials and Theory of Structures.	8vo,	7 50
Burr's Elasticity and Resistance of the Materials of Engineering. 6th Edition. Reset.	8vo,	7 50
Church's Mechanics of Engineering.	8vo,	6 00
* Greene's Structural Mechanics	8vo,	2 50
Johnson's Materials of Construction.	8vo,	6 00
Keep's Cast Iron.	8vo,	2 50
Lanza's Applied Mechanics.	8vo,	7 50
Martens's Handbook on Testing Materials. (Henning.)	8vo,	7 50
Maurer's Technical Mechanics.	8vo,	4 00
Merriman's Mechanics of Materials.	8vo,	5 00
Strength of Materials	12mo,	1 00
Metcalf's Steel. A manual for Steel-users.	12mo,	2 00
Sabin's Industrial and Artistic Technology of Paints and Varnish.	8vo,	3 00
Smith's Materials of Machines.	12mo,	1 00
Thurston's Materials of Engineering.	3 vols, 8vo,	8 00
Part II. Iron and Steel.	8vo,	3 50
Part III. A Treatise on Brasses, Bronzes, and Other Alloys and their Constituents.	8vo,	2 50
Text-book of the Materials of Construction.	8vo,	5 00
Wood's (De V.) Treatise on the Resistance of Materials and an Appendix on the Preservation of Timber.	8vo,	2 00

Wood's (De V.) Elements of Analytical Mechanics.	8vo,	3 00
Wood's (M. P.) Rustless Coatings: Corrosion and Electrolysis of Iron and Steel.	8vo,	4 00

STEAM-ENGINES AND BOILERS.

Berry's Temperature-entropy Diagram.	12mo,	1 25
Carnot's Reflections on the Motive Power of Heat. (Thurston.)	12mo,	1 50
Dawson's "Engineering" and Electric Traction Pocket-book.	16mo, mor.,	5 00
Ford's Boiler Making for Boiler Makers.	18mo,	1 00
Goss's Locomotive Sparks.	8vo,	2 00
Hemenway's Indicator Practice and Steam-engine Economy.	12mo,	2 00
Hutton's Mechanical Engineering of Power Plants.	8vo,	5 00
Heat and Heat-engines.	8vo,	5 00
Kent's Steam boiler Economy.	8vo,	4 00
Kneass's Practice and Theory of the Injector.	8vo,	1 50
MacCord's Slide-valves.	8vo,	2 00
Meyer's Modern Locomotive Construction.	4to,	10 00
Peabody's Manual of the Steam-engine Indicator.	12mo,	1 50
Tables of the Properties of Saturated Steam and Other Vapors	8vo,	1 00
Thermodynamics of the Steam-engine and Other Heat-engines.	8vo,	5 00
Valve-gears for Steam-engines.	8vo,	2 50
Peabody and Miller's Steam-boilers.	8vo,	4 00
Pray's Twenty Years with the Indicator.	Large 8vo,	2 50
Pupin's Thermodynamics of Reversible Cycles in Gases and Saturated Vapors. (Osterberg.)	12mo,	1 25
Reagan's Locomotives: Simple Compound, and Electric.	12mo,	2 50
Rontgen's Principles of Thermodynamics. (Du Bois.)	8vo,	5 00
Sinclair's Locomotive Engine Running and Management.	12mo,	2 00
Smart's Handbook of Engineering Laboratory Practice.	12mo,	2 50
Snow's Steam-boiler Practice.	8vo,	3 00
Spangler's Valve-gears.	8vo,	2 50
Notes on Thermodynamics.	12mo,	1 00
Spangler, Greene, and Marshall's Elements of Steam-engineing	8vo,	3 00
Thurston's Handy Tables.	8vo,	1 50
Manual of the Steam-engine.	2 vols., 8vo,	10 00
Part I. History, Structure, and Theory.	8vo,	6 00
Part II. Design, Construction, and Operation.	8vo,	6 00
Handbook of Engine and Boiler Trials, and the Use of the Indicator and the Prony Brake.	8vo,	5 00
Stationary Steam-engines.	8vo,	2 50
Steam-boiler Explosions in Theory and in Practice	12mo,	1 50
Manual of Steam-boilers, their Designs, Construction, and Operation.	8vo,	5 00
Weisbach's Heat, Steam, and Steam-engines. (Du Bois.)	8vo,	5 00
Whitham's Steam-engine Design.	8vo,	5 00
Wilson's Treatise on Steam-boilers. (Flather.)	16mo,	2 50
Wood's Thermodynamics, Heat Motors, and Refrigerating Machines.	8vo,	4 00

MECHANICS AND MACHINERY.

Barr's Kinematics of Machinery.	8vo,	2 50
* Bovey's Strength of Materials and Theory of Structures	8vo,	7 50
Chase's The Art of Pattern-making.	12mo,	2 50
Church's Mechanics of Engineering.	8vo,	6 00
Notes and Examples in Mechanics.	8vo,	2 00
Compton's First Lessons in Metal-working.	12mo,	1 50
Compton and De Groodt's The Speed Lathe.	12mo,	1 50

Cromwell's Treatise on Toothed Gearing.....	12mo,	1 50
Treatise on Belts and Pulleys.....	12mo,	7 50
Dana's Text-book of Elementary Mechanics for Colleges and Schools.....	12mo,	1 50
Dingey's Machinery Pattern Making.....	12mo,	2 00
Dredge's Record of the Transportation Exhibits Building of the World's Columbian Exposition of 1893.....	4to half morocco,	5 00
Du Bois's Elementary Principles of Mechanics:		
Vol. I. Kinematics.....	8vo,	3 50
Vol. II. Statics.....	8vo,	4 00
Mechanics of Engineering. Vol. I.....	Small 4to,	7 50
Vol. II.....	Small 4to,	10 00
Durley's Kinematics of Machines.....	8vo,	4 00
Fitzgerald's Boston Machinist.....	16mo,	1 00
Flather's Dynamometers, and the Measurement of Power.....	12mo,	3 00
Rope Driving.....	12mo,	2 00
Goss's Locomotive Sparks.....	8vo,	2 00
* Greene's Structural Mechanics.....	8vo,	2 50
Hall's Car Lubrication.....	12mo,	1 00
Holly's Art of Saw Filing.....	18mo,	75
James's Kinematics of a Point and the Rational Mechanics of a Particle.....	Small 8vo,	2 00
* Johnson's (W. W.) Theoretical Mechanics.....	12mo,	3 00
Johnson's (L. J.) Statics by Graphic and Algebraic Methods.....	8vo,	2 00
Jones's Machine Design:		
Part I. Kinematics of Machinery.....	8vo,	1 50
Part II. Form, Strength, and Proportions of Parts.....	8vo,	3 00
Kerr's Power and Power Transmission.....	8vo,	2 00
Lanza's Applied Mechanics.....	8vo,	7 50
Leonard's Machine Shop, Tools, and Methods.....	8vo,	4 00
* Lorenz's Modern Refrigerating Machinery. (Pope, Haven, and Dean.).....	8vo,	4 00
MacCord's Kinematics; or, Practical Mechanism.....	8vo,	5 00
Velocity Diagrams.....	8vo,	1 50
Maurer's Technical Mechanics.....	8vo,	4 00
Merriman's Mechanics of Materials.....	8vo,	5 00
* Elements of Mechanics.....	12mo,	1 00
* Michie's Elements of Analytical Mechanics.....	8vo,	4 00
Reagan's Locomotives: Simple, Compound, and Electric.....	12mo,	2 50
Reid's Course in Mechanical Drawing.....	8vo,	2 00
Text-book of Mechanical Drawing and Elementary Machine Design.....	8vo,	3 00
Richards's Compressed Air.....	12mo,	1 50
Robinson's Principles of Mechanism.....	8vo,	3 00
Ryan, Norris, and Hoxie's Electrical Machinery. Vol. I.....	8vo,	2 50
Schwamb and Merrill's Elements of Mechanism.....	8vo,	3 00
Sinclair's Locomotive-engine Running and Management.....	12mo,	2 00
Smith's (O.) Press-working of Metals.....	8vo,	3 00
Smith's (A. W.) Materials of Machines.....	12mo,	1 00
Smith (A. W.) and Marx's Machine Design.....	8vo,	3 00
Spangler, Greene, and Marshall's Elements of Steam-engineering.....	8vo,	3 00
Thurston's Treatise on Friction and Lost Work in Machinery and Mill Work.....	8vo,	3 00
Animal as a Machine and Prime Motor, and the Laws of Energetics.....	12mo,	1 00
Warren's Elements of Machine Construction and Drawing.....	8vo,	7 50
Weisbach's Kinematics and Power of Transmission. (Herrmann—Klein.).....	8vo,	5 00
Machinery of Transmission and Governors. (Herrmann—Klein.).....	8vo,	5 00
Wood's Elements of Analytical Mechanics.....	8vo,	3 00
Principles of Elementary Mechanics.....	12mo,	1 25
Turbines.....	8vo,	2 50
The World's Columbian Exposition of 1893.....	4to,	1 00

METALLURGY.

Egleston's Metallurgy of Silver, Gold, and Mercury:		
Vol. I. Silver.	8vo,	7 50
Vol. II. Gold and Mercury.	8vo,	7 50
** Iles's Lead-smelting. (Postage 9 cents additional.)	12mo,	2 50
Keep's Cast Iron.	8vo,	2 50
Kunhardt's Practice of Ore Dressing in Europe.	8vo,	1 50
Le Chatelier's High-temperature Measurements. (Boudouard—Burgess.)	12mo,	3 00
Metcalf's Steel. A Manual for Steel-users.	12mo,	2 00
Minet's Production of Aluminum and its Industrial Use. (Waldo.)	12mo,	2 50
Robine and Lenglen's Cyanide Industry. (Le Clerc.)	8vo,	
Smith's Materials of Machines.	12mo,	1 00
Thurston's Materials of Engineering. In Three Parts.	8vo,	8 00
Part II. Iron and Steel.	8vo,	3 50
Part III. A Treatise on Brasses, Bronzes, and Other Alloys and their Constituents.	8vo,	2 50
Ulke's Modern Electrolytic Copper Refining.	8vo,	3 00

MINERALOGY.

Barringer's Description of Minerals of Commercial Value. Oblong, morocco.	2 50
Boyd's Resources of Southwest Virginia.	8vo, 3 00
Map of Southwest Virginia.	Pocket-book form. 2 00
Brush's Manual of Determinative Mineralogy. (Penfield.)	8vo, 4 00
Chester's Catalogue of Minerals.	8vo, paper, 1 00
	Cloth, 1 25
Dictionary of the Names of Minerals.	8vo, 3 50
Dana's System of Mineralogy.	Large 8vo, half leather, 12 50
First Appendix to Dana's New "System of Mineralogy."	Large 8vo, 1 00
Text-book of Mineralogy.	8vo, 4 00
Minerals and How to Study Them.	12mo. 1 50
Catalogue of American Localities of Minerals.	Large 8vo, 1 00
Manual of Mineralogy and Petrography.	12mo, 2 00
Douglas's Untechnical Addresses on Technical Subjects.	12mo, 1 00
Eakle's Mineral Tables.	8vo, 1 25
Egleston's Catalogue of Minerals and Synonyms.	8vo, 2 50
Hussak's The Determination of Rock-forming Minerals. (Smith.)	Small 8vo, 2 00
Merrill's Non-metallic Minerals: Their Occurrence and Uses.	8vo, 4 00
* Penfield's Notes on Determinative Mineralogy and Record of Mineral Tests. 8vo, paper,	50
Rosenbusch's Microscopical Physiography of the Rock-making Minerals. (Iddings.)	8vo, 5 00
* Tillman's Text-book of Important Minerals and Rocks.	8vo, 2 00

MINING.

Beard's Ventilation of Mines.	12mo, 2 50
Boyd's Resources of Southwest Virginia.	8vo, 3 00
Map of Southwest Virginia.	Pocket-book form 2 00
Douglas's Untechnical Addresses on Technical Subjects.	12mo, 1 00
* Drinker's Tunneling, Explosive Compounds, and Rock Drills. .4to, hf. mor.,	25 00
Eissler's Modern High Explosives.	8vo. 4 00

Fowler's Sewage Works Analyses.....	12mo,	2 00
Goodyear's Coal-mines of the Western Coast of the United States.....	12mo,	2 50
Ihlseng's Manual of Mining.....	8vo,	5 00
** Jles's Lead-smelting. (Postage 9c. additional.).....	12mo,	2 50
Kunhardt's Practice of Ore Dressing in Europe.....	8vo,	1 50
O'Driscoll's Notes on the Treatment of Gold Ores.....	8vo,	2 00
Robine and Lenglen's Cyanide Industry. (Le Clerc.).....	8vo,	
* Walke's Lectures on Explosives.....	8vo,	4 00
Wilson's Cyanide Processes.....	12mo,	1 50
Chlorination Process.....	12mo,	1 50
Hydraulic and Placer Mining.....	12mo,	2 00
Treatise on Practical and Theoretical Mine Ventilation.....	12mo,	1 25

SANITARY SCIENCE.

Bashore's Sanitation of a Country House.....	12mo,	1 00
Folwell's Sewerage. (Designing, Construction, and Maintenance.).....	8vo,	3 00
Water-supply Engineering.....	8vo,	4 00
Fuertes's Water and Public Health.....	12mo,	1 50
Water-filtration Works.....	12mo,	2 50
Gerhard's Guide to Sanitary House-inspection.....	16mo,	1 00
Goodrich's Economic Disposal of Town's Refuse.....	Demy 8vo,	3 50
Hazen's Filtration of Public Water-supplies.....	8vo,	3 00
Leach's The Inspection and Analysis of Food with Special Reference to State Control.....	8vo,	7 50
Mason's Water-supply. (Considered principally from a Sanitary Standpoint).....	8vo,	4 00
Examination of Water. (Chemical and Bacteriological.).....	12mo,	1 25
Ogden's Sewer Design.....	12mo,	2 00
Prescott and Winslow's Elements of Water Bacteriology, with Special Reference to Sanitary Water Analysis.....	12mo,	1 25
* Price's Handbook on Sanitation.....	12mo,	1 50
Richards's Cost of Food. A Study in Dietaries.....	12mo,	1 00
Cost of Living as Modified by Sanitary Science.....	12mo,	1 00
Richards and Woodman's Air, Water, and Food from a Sanitary Standpoint.....	8vo,	2 00
* Richards and Williams's The Dietary Computer.....	8vo,	1 50
Rideal's Sewage and Bacterial Purification of Sewage.....	8vo,	3 50
Turneure and Russell's Public Water-supplies.....	8vo,	5 00
Von Behring's Suppression of Tuberculosis. (Bolduan.).....	12mo,	1 00
Whipple's Microscopy of Drinking-water.....	8vo,	3 50
Winton's Microscopy of Vegetable Foods.....	8vo,	7 50
Woodhull's Notes on Military Hygiene.....	16mo,	1 50

MISCELLANEOUS.

De Fursac's Manual of Psychiatry. (Rosanoff and Collins.)....	Large 12mo,	2 50
Emmons's Geological Guide-book of the Rocky Mountain Excursion of the International Congress of Geologists.....	Large 8vo,	1 50
Ferrel's Popular Treatise on the Winds.....	8vo,	4 00
Haines's American Railway Management.....	12mo,	2 50
Mott's Fallacy of the Present Theory of Sound.....	16mo,	1 00
Ricketts's History of Rensselaer Polytechnic Institute, 1824-1894..	Small 8vo,	3 00
Rostoski's Serum Diagnosis. (Bolduan.).....	12mo,	1 00
Rotherham's Emphasized New Testament.....	Large 8vo,	2 00

Steel's Treatise on the Diseases of the Dog.	8vo.	3	50
The World's Columbian Exposition of 1893	4to,	1	00
Von Behring's Suppression of Tuberculosis. (Bolduan.)	12mo,	1	00
Winslow's Elements of Applied Microscopy	12mo,	1	50
Worcester and Atkinson. Small Hospitals, Establishment and Maintenance; Suggestions for Hospital Architecture: Plans for Small Hospital.	12mo,	1	25

HEBREW AND CHALDEE TEXT-BOOKS.

Green's Elementary Hebrew Grammar	12mo,	1	25
Hebrew Chrestomathy.	8vo,	2	00
Gesenius's Hebrew and Chaldee Lexicon to the Old Testament Scriptures. (Tregelles.)	Small 4to, half morocco,	5	00
Letteris's Hebrew Bible.	8vo,	2	25



$S_1 \epsilon$
 $\overline{p} \rightarrow 0$ m

14 DAY USE
RETURN TO DESK FROM WHICH BORROWED
LOAN DEPT.

This book is due on the last date stamped below,
or on the date to which renewed. Renewals only:
Tel. No. 642-3405
Renewals may be made 4 days prior to date due.
Renewed books are subject to immediate recall.

Stanford

INTER-LIBRARY
LOAN

OCT 26 1971

DEC 30 1975

SANTA BARBARA

INTERLIBRARY LOAN RECEIVED BY

REC. CIR. JAN 30 '76

APR 26 1986

CIRCULATION DEPT.

MAY 12 1987

SEP 08 '91

SEP 5 1982


RECEIVED BY AUTO DISC AUG 12 '91

AUG 11 1982

CIRCULATION DEPT.
LD 000 000 000 000 000 000
(P2001s10)476-A-32

General Library
University of California
Berkeley

GENERAL LIBRARY - U.C. BERKELEY



8000905427

Q4257
G7549

236367

